

## Shell Matrices and Fermion Vertices-Predicational Anteriority and Character Constitution Thereof

<sup>1</sup>Dr. K. N. Prasanna Kumar, <sup>2</sup>Prof. B. S. Kiranagi, <sup>3</sup>Prof. C. S. Bagewadi

**Abstract:** In quantum physics, in order to quantize a gauge theory, like for example Yang-Mills theory, Chern-Simons or BF model, one method is to perform a gauge fixing. This is done in the BRST and Batalin-Vilkovisky formulation. Another is to factor out the symmetry by dispensing with vector potentials altogether (they're not physically observable anyway) and work directly with Wilson loops, Wilson lines contracted with other charged fields at its endpoints and spin networks Presently renormalization prescriptions of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix have been investigated by many authors like Yong Zhou. Based on one prescription which is formulated by comparing with the fictitious case of no mixing of quark generations, they have proposed the substantive limits, singular pauses, general rests, logical attributes and real relations therefore a new prescription intermodal manifestation which can make the physical amplitude involving quark's mixing gauge independent and ultraviolet finite. Compared with the previous prescriptions this prescription is very simple and suitable for actual calculations. Through analytical calculations we also give a strong Proof for the important hypothesis that in order to keep the CKM matrix gauge independent the unitarity of the CKM matrix must be preserved. Mass-shell renormalization of fermion mixing matrices have also been delineated and investigated upon by K.-P.O Diener, B.A Kniehl wherein they consider favorable extensions of the standard model (SM) where the lepton sector contains Majorana neutrinos with vanishing left-handed mass terms, thus allowing for the see-saw mechanism to operate, and propose physical on-mass-shell (OS) renormalization conditions for the lepton mixing matrices that comply with ultraviolet finiteness, gauge-parameter independence, and (pseudo)unitarity This is an important result that motivated us to draw up the consolidation of some of the most important variables in Fermion and graviton vertices.. A crucial feature is that the texture zero in the neutrino mass matrix is preserved by renormalization, which is not automatically the case for possible generalizations of existing renormalization prescriptions for the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix in the SM. Our renormalization prescription also applies to the special case of the SM and leads to a physical OS definition of the renormalized CKM matrix. A consummate and link model is built for the variables like gravity, matter field, virtual photons and other important variables. Nevertheless the stormy petrel Neutrino seems to rule the roost with its own disnormative prescriptions for itself. Rich IN ITS twists and turns, the Model seems to offer a parade of variables bent on aggrandizement agenda.

### I. Introduction:

#### Ward-Takahashi identity

In quantum field theory, a Ward-Takahashi identity is an identity between correlation functions that follows from the global or gauged symmetries of the theory, and which remains valid after renormalization.

The Ward-Takahashi identity of quantum electrodynamics was originally used by John Clive Ward and Yasushi Takahashi to relate the wave function renormalization of the electron to its vertex renormalization factor  $F_1(0)$ , guaranteeing the cancellation of the ultraviolet divergence to all orders of perturbation theory. Later uses include the extension of the proof of Goldstone's theorem to all orders of perturbation theory.

The Ward-Takahashi identity is a quantum version of the classical Noether's theorem, and any symmetry in a quantum field theory can lead to an equation of motion for correlation functions..

The Ward-Takahashi identity applies to correlation functions in momentum space, which do not necessarily have all their external momenta on-shell. Let

$$\mathcal{M}(k; p_1 \cdots p_n; q_1 \cdots q_n) = \epsilon_\mu(k) \mathcal{M}^\mu(k; p_1 \cdots p_n; q_1 \cdots q_n)$$

be a QED correlation function involving an external photon with momentum  $k$  (where  $\epsilon_\mu(k)$  is the polarization vector of the photon),  $n$  initial-state electrons with momenta  $p_1 \cdots p_n$ , and  $n$  final-state electrons with momenta  $q_1 \cdots q_n$ .

Also define  $\mathcal{M}_0$  to be the simpler amplitude that is **obtained by removing** the photon with momentum  $k$  from original amplitude. Then the Ward-Takahashi identity reads

$$k_\mu \mathcal{M}^\mu(k; p_1 \cdots p_n; q_1 \cdots q_n) = -e \sum_i [\mathcal{M}_0(p_1 \cdots p_n; q_1 \cdots (q_i - k) \cdots q_n) - \mathcal{M}_0(p_1 \cdots (p_i + k) \cdots p_n; q_1 \cdots q_n)]$$

where  $-e$  is the charge of the electron. Note that if  $\mathcal{M}$  has its external electrons on-shell, then the amplitudes on the right-hand side of this identity each had one external particle off-shell, and therefore they do not contribute to S-matrix elements.

#### The Ward identity

The Ward identity is a specialization of the Ward-Takahashi identity to S-matrix elements, which describe physically possible scattering processes and thus have all their external particles on-shell. Again let  $\mathcal{M}(k) = \epsilon_\mu(k) \mathcal{M}^\mu(k)$

be the amplitude for some QED process involving an external photon with momentum  $k$ , where  $\epsilon_\mu(k)$  is the polarization vector of the photon. Then the Ward identity reads:

$$k_\mu \mathcal{M}^\mu(k) = 0$$

Physically, what this identity means is the longitudinal polarization of the photon which arises in the  $\xi$  gauge is unphysical and disappears from the S-matrix.

## II. Some Reviews:

Flavor Changing Fermion-Graviton Vertices (SEE FOR DETAILS G. Degrassi, E. Gabrielli, L. Trentadue-emphasis mine) Authors study the flavor-changing quark-graviton vertex that is induced at the one-loop level when gravitational interactions are coupled to the standard model. Because of the conservation of the energy-momentum tensor the corresponding form factors turn out to be finite and gauge-invariant. Analytical expressions of the form factors are provided at leading order in the external masses. Authors show that flavor-changing interactions in gravity are local if the graviton is strictly massless while if the graviton has a small mass long-range interactions inducing a flavor-changing contribution in the Newton potential appear. Flavor-changing processes with massive spin-2 particles are also briefly discussed in the paper. . These results can be generalized to the case of the lepton-graviton coupling.

Examples of its use include constraining the tensor structure of the vacuum polarization and of the electron vertex function in QED. Gauge dependence of the on-shell renormalized mixing matrices WAS STUDIED BY Youichi Yamada It was recently pointed out that the on-shell renormalization of the CabibboKobayashi-Maskawa (CKM) matrix in the method by Denner and Sack causes a gauge parameter dependence of the amplitudes. Authors analyze the gauge dependence of the on-shell renormalization of the mixing matrices both for fermions and scalars in general cases, at the one-loop level. It is also shown that this gauge dependence can be avoided by fixing the counterterm for the mixing matrices in terms of the off-diagonal wave function corrections for fermions and scalars after a rearrangement, in a similar manner to the pinch technique for gauge bosons. Particles in the same representation under unbroken symmetries can mix with each other. The neutral gauge bosons, quarks, and massive neutrinos in the Standard Model (SM) are well-known examples. New particles in extensions of the Standard Model also show the mixings. For example, in the minimal supersymmetric (SUSY) standard model (MSSM), a very promising extension, super partners of most SM particles show the mixing .The mixing of particles is expressed in terms of the mixing matrix, which represents the relations between the gauge eigenstates and the mass eigenstates of the particles. The mixing matrices always appear at the couplings of these particles in the mass eigenbasis. Because of the fact that mass eigenstates at the tree-level mix with each other by radiative corrections, (it calls for) the mixing matrices have to be renormalized to obtain ultraviolet (UV) finite amplitudes. Denner and Sack have proposed a simple scheme to renormalize the mixing matrix of Dirac fermions at the one-loop level, which is usually called the on-shell renormalization scheme. They have required the counterterm for the renormalized mixing matrix to completely absorb the anti-Hermitian part of the wave function correction  $\delta Z_{ij}$  for the external on-shell fields\*

## III. GRAVITY AND MATTER FIELDS:

### MODULE NUMBERED ONE NOTATION :

- $G_{13}$  : CATEGORY ONE OF GRAVITY
- $G_{14}$  : CATEGORY TWO OF GRAVITY
- $G_{15}$  : CATEGORY THREE OF GRAVITY
- $T_{13}$  : CATEGORY ONE OF MATTER FIELDS
- $T_{14}$  : CATEGORY TWO OF MATTER FIELDS
- $T_{15}$  : CATEGORY THREE OF MATTER FIELDS

GRAVITON FIELD AND CONSERVED MATTER ENERGY MOMENTUM TENSOR(LIKE IN A BANK THE RULE THAT ASSETS AND LIABILITIES ARE EQUIVALENT IS APPLIED TO THE INDIVIDUAL SYSTEMS, THE CONSERVATION OF ENERGY MOMENTUM TENSOR IS APPLICABLE TO VARIOUS SYSTEMS AND THE CLASSIFICATION IS BASED ON THE CHARACTERISTICS OF THE SYSTEMS TO WHICH THE CONSERVATION PRINCIPLE IS APPLIED):

### MODULE NUMBERED TWO:

- $G_{16}$  : CATEGORY ONE OF GRAVITON FIELD
- $G_{17}$  : CATEGORY TWO OF GRAVITON FIELD
- $G_{18}$  : CATEGORY THREE OF GRAVITON FIELD
- $T_{16}$  : CATEGORY ONE OF CONSERVED MATTER-ENERGY-MOMENTUM TENSOR(WE ARE HERE SPEAKING OF SYSTEMS TO WHICH IT IS APPLICABLE. PLEASE THE BANK EXAMPLE GIVEN ABOVE)
- $T_{17}$  : CATEGORY TWO OF CONSERVED MATTER-ENERGY-MOMENTUM TENSOR
- $T_{18}$  : CATEGORY THREE OF CONSERVED MATTER-ENERGY-MOMENTUM TENSOR

### VIRTUAL PHOTONS AND GRAVITON PHOTON VERTEX:

#### MODULE NUMBERED THREE:

$G_{20}$  : CATEGORY ONE OF VIRTUAL PHOTONS(WE HERE SPEAK OF THE CHARACTERISED SYSTEMS FOR WHICH QUANTUM GAUGE THEORY IS APPLICABLE)

$G_{21}$  :CATEGORY TWO OF VIRTUAL PHOTONS

$G_{22}$  : CATEGORY THREE OF VIRTUAL PHOTONS

$T_{20}$  : CATEGORY ONE OF GRAVITON ELECTRON VERTEX

$T_{21}$  :CATEGORY TWO OF GRAVITON ELECTRON VERTEX

$T_{22}$  : CATEGORY THREE OF GRAVITON ELECTRON VERTEX

**QUANTUM FIELD THEORY(AGAIN,PARAMETRICIZED SYSTEMS TO WHICH QFT COULD BE APPLIED IS TAKEN IN TO CONSIDERATION AND RENORMALIZATION THEORY(BASED ON CERTAIN VARIABLES OF THE SYSTEM WHICH CONSEQUENTIALLY CLSSIFIABLE ON PARAMETERS)**  
**: MODULE NUMBERED FOUR:**

$G_{24}$  : CATEGORY ONE OF QUANTUM FIELD THEORY(EVALUATIVE PARAMETRICIZATION OF SITUATIONAL ORIENTATIONS AND ESSENTIAL COGNITIVE ORIENTATION AND CHOICE VARIABLES OF THE SYSTEM TO WHICH QFT IS APPLICABLE)

$G_{25}$  : CATEGORY TWO OF QUANTUM FIELD THEORY

$G_{26}$  : CATEGORY THREE OF QUANTUM FIELD THEORY

$T_{24}$  :CATEGORY ONE OF RENORMALIZATION THEORY

$T_{25}$  :CATEGORY TWO OF RENORMALIZATION THEORY(SYSTEMIC INSTRUMENTAL CHARACTERISATIONS AND ACTION ORIENTATIONS AND FUYNCTIONAL IMPERATIVES OF CHANGE MANIFESTED THEREIN )

$T_{26}$  : CATEGORY THREE OF QUANTUM FIELD THEORY

**VIRTUAL ELECTRONS AND GRAVITON PHOTON VERTEX**  
**MODULE NUMBERED FIVE:**

$G_{28}$  : CATEGORY ONE OF VIRTUAL ELECTRONS

$G_{29}$  : CATEGORY TWO OF VIRTUAL ELECTRONS

$G_{30}$  :CATEGORY THREE OF VIRTUAL ELECTRONS

$T_{28}$  :CATEGORY ONE OF GRAVITON PHOTON VERTEX

$T_{29}$  :CATEGORY TWO OF GRAVITON PHOTON VERTEX

$T_{30}$  :CATEGORY THREE OF GRAVITON PHOTON VERTEX

**QUANTUM CORRECTIONS TO ON SHELL MATRIX (VIRTUAKL GRAVITONS ARE NOT INCLUDED IN THE LOOPS) AND WARD IDENTITIES FROM MATTER ENERGY MOMENTUM CONSERVATION(LOT OF SYSTEMS CONSERVE THE MASS ENERGY AND THE CLASSIFICATION IS BASED ON THE PARAMETRICIZATION OF THE SYSTEMS)**  
**MODULE NUMBERED SIX:**

$G_{32}$  : CATEGORY ONE OF QUANTUM CORRECTION TO SHELL MATRIX

$G_{33}$  : CATEGORY TWO OF QUANTUM CORRECTIONS TO SHELL MATRIX

$G_{34}$  : CATEGORY THREE OF QUANTUM CORRECTIONS TO SHELL MATRIX

$T_{32}$  : CATEGORY ONE OF WARD IDENTITIES FROM MASS-ENERGY-MOMENTUM CONSERVATION(AGAIN WE RECAPITUALTE THE BANK EXAMPLES THERE ARE MILLIONS OF SYSTEMS FOR WHICH THE CONSERVATION HOLDS AND WE ARE CLASSIFYING THE SYSTEMS AND WARD IDENTITIES THEREOF)

$T_{33}$  : CATEGORY TWO OF WARD IDENTITIES

$T_{34}$  : CATEGORY THREE OF WARD IDENTITIES

**CHARGED WEAK CURRENTS AND ONE LOOP FLAVOUR CHANGING NEUTRAL CURRENTS(FCNC) IN THE FERMION PORTFOLIO:**  
**MODULE NUMBERED SEVEN**

$G_{36}$  : CATEGORY ONE OF CHARGED WEAK CURRENTS

$G_{37}$  : CATEGORY TWO OF CHARGED WEAK CURRENTS

$G_{38}$  : CATEGORY THREE OF CHARGED WEAK CURRENTS (ENERGY EXCITATION OF THE VACUUM AND CONCOMITANT GENERATION OF ENERGY DIFFERENTIAL-TIME LAG OR INSTANTANEOUSNESSMIGHT EXISTS WHEREBY ACCENTUATION AND ATTRITIONS MODEL MAY ASSUME ZERO POSITIONS IS AN EXAMPLE)

$T_{36}$  : CATEGORY ONE OF FCNC IN THE FERMIONS SECTOR

$T_{37}$  : CATEGORY TWO OF FCNC IN THE FERMIONS SECTOR

$T_{38}$  : CATEGORY THREE OF FCNC IN THE FERMIONS SECTOR

$$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)} \quad (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)} \quad (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$$

$$(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$$

$$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$$

$$(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$$

$$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$$

$$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)} \quad (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$$

$$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

are Dissipation coefficients\*

**GRAVITY AND MATTER FIELDS:**  
**MODULE NUMBERED ONE**

The differential system of this model is now (Module Numbered one)\*1

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad *2$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad *3$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad *4$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad *5$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad *6$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad *7$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \quad *8$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor} \quad *$$

**GRAVITON FIELD AND CONSERVED MATTER ENERGY MOMENTUM TENSOR(LIKE IN A BANK THE RULE THAT ASSETS AND LIABILITIES ARE EQUIVALENT IS APPLIED TO THE INDIVIDUAL SYSTEMS, THE CONSERVATION OF ENERGY MOMENTUM TENSOR IS APPLICABLE TO VARIOUS SYSTEMS AND THE CLASSIFICATION IS BASED ON THE CHARACTERISTICS OF THE SYSTEMS TO WHICH THE CONSERVATION PRINCIPLE IS APPLIED):**

**MODULE NUMBERED TWO:**

∴

The differential system of this model is now (Module numbered two)\*9

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad *10$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad *11$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad *12$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad *13$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad *14$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad *15$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \quad *16$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \quad *17$$

**VIRTUAL PHOTONS AND GRAVITON PHOTON VERTEX:**  
**MODULE NUMBERED THREE**

The differential system of this model is now (Module numbered three)\*18

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad *19$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad *20$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad *21$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad *22$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad *23$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad *24$$

$$+(a_{20}'')^{(3)}(T_{21}, t) = \text{First augmentation factor}^*$$

$$-(b_{20}'')^{(3)}(G_{23}, t) = \text{First detritions factor} *25$$

**QUANTUM FIELD THEORY (AGAIN, PARAMETRIZED SYSTEMS TO WHICH QFT COULD BE APPLIED IS TAKEN IN TO CONSIDERATION AND RENORMALIZATION THEORY (BASED ON CERTAIN VARIABLES OF THE SYSTEM WHICH CONSEQUENTIALLY CLASSIFIABLE ON PARAMETERS)**

**: MODULE NUMBERED FOUR**

The differential system of this model is now (Module numbered Four)\*26

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a_{24}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)]G_{24} *27$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a_{25}')^{(4)} + (a_{25}'')^{(4)}(T_{25}, t)]G_{25} *28$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a_{26}')^{(4)} + (a_{26}'')^{(4)}(T_{25}, t)]G_{26} *29$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b_{24}')^{(4)} - (b_{24}'')^{(4)}((G_{27}, t))]T_{24} *30$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b_{25}')^{(4)} - (b_{25}'')^{(4)}((G_{27}, t))]T_{25} *31$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b_{26}')^{(4)} - (b_{26}'')^{(4)}((G_{27}, t))]T_{26} *32$$

$$+(a_{24}'')^{(4)}(T_{25}, t) = \text{First augmentation factor} *33$$

$$-(b_{24}'')^{(4)}((G_{27}, t)) = \text{First detritions factor} *34$$

**VIRTUAL ELECTRONS AND GRAVITON PHOTON VERTEX**

**MODULE NUMBERED FIVE**

The differential system of this model is now (Module number five)\*35

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a_{28}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)]G_{28} *36$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a_{29}')^{(5)} + (a_{29}'')^{(5)}(T_{29}, t)]G_{29} *37$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a_{30}')^{(5)} + (a_{30}'')^{(5)}(T_{29}, t)]G_{30} *38$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b_{28}')^{(5)} - (b_{28}'')^{(5)}((G_{31}, t))]T_{28} *39$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b_{29}')^{(5)} - (b_{29}'')^{(5)}((G_{31}, t))]T_{29} *40$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b_{30}')^{(5)} - (b_{30}'')^{(5)}((G_{31}, t))]T_{30} *41$$

$$+(a_{28}'')^{(5)}(T_{29}, t) = \text{First augmentation factor} *42$$

$$-(b_{28}'')^{(5)}((G_{31}, t)) = \text{First detritions factor} *43$$

**QUANTUM CORRECTIONS TO ON SHELL MATRIX (VIRTUAKL GRAVITONS ARE NOT INCLUDED IN THE LOOPS) AND WARD IDENTITIES FROM MATTER ENERGY MOMENTUM CONSERVATION (LOT OF SYSTEMS CONSERVE THE MASS ENERGY AND THE CLASSIFICATION IS BASED ON THE PARAMETRIZATION OF THE SYSTEMS)**

**MODULE NUMBERED SIX:**

The differential system of this model is now (Module numbered Six)\*44

45

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a_{32}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)]G_{32} *46$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a_{33}')^{(6)} + (a_{33}'')^{(6)}(T_{33}, t)]G_{33} *47$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a_{34}')^{(6)} + (a_{34}'')^{(6)}(T_{33}, t)]G_{34} *48$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b_{32}')^{(6)} - (b_{32}'')^{(6)}((G_{35}, t))]T_{32} *49$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b_{33}')^{(6)} - (b_{33}'')^{(6)}((G_{35}, t))]T_{33} *50$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b_{34}')^{(6)} - (b_{34}'')^{(6)}((G_{35}, t))]T_{34} *51$$

$$+(a_{32}'')^{(6)}(T_{33}, t) = \text{First augmentation factor} *52$$

**CHARGED WEAK CURRENTS AND ONE LOOP FLAVOUR CHANGING NEUTRAL CURRENTS (FCNC) IN THE FERMION PORTFOLIO:**

**MODULE NUMBERED SEVEN**

The differential system of this model is now (SEVENTH MODULE)\*53

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a_{36}')^{(7)} + (a_{36}'')^{(7)}(T_{37}, t)]G_{36} *54$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a_{37}')^{(7)} + (a_{37}'')^{(7)}(T_{37}, t)]G_{37} *55$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a_{38}')^{(7)} + (a_{38}'')^{(7)}(T_{37}, t)]G_{38} *56$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad *57$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad *58$$

59

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad *60$$

$$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \quad *61$$

$$-(b''_{36})^{(7)}((G_{39}), t) = \text{First detritions factor}$$

**FIRST MODULE CONCATENATION:**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{c} (a'_{13})^{(1)} \boxed{+(a''_{13})^{(1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7)}(T_{37}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{c} (a'_{14})^{(1)} \boxed{+(a''_{14})^{(1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7)}(T_{37}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{c} (a'_{15})^{(1)} \boxed{+(a''_{15})^{(1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7)}(T_{37}, t)} \end{array} \right] G_{15}$$

Where  $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{36})^{(7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7)}(T_{37}, t)}$  ARE SEVENTH AUGMENTATION COEFFICIENTS

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{c} (b'_{13})^{(1)} \boxed{-(b''_{16})^{(1)}(G, t)} \boxed{-(b''_{36})^{(7)}(G_{39}, t)} \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7)}(G_{39}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{c} (b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7)}(G_{39}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{c} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7)}(G_{39}, t)} \end{array} \right] T_{15}$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$  are fifth detritions coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$  are sixth detritions coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7)}(G_{39}, t)$  ARE SEVENTH DETRITION COEFFICIENTS

\*62

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{ccc} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} & \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{15} \quad *63$$

Where  $-(b''_{13})^{(1)}(G, t)$ ,  $-(b''_{14})^{(1)}(G, t)$ ,  $-(b''_{15})^{(1)}(G, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2)}(G_{19}, t)$  are second detritions coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3)}(G_{23}, t)$  are third detritions coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$  are fourth detritions coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$  are fifth detritions coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6)}(G_{35}, t)$  are sixth detritions coefficients for category 1, 2 and 3 \*64

**SECOND MODULE CONCATENATION: \*65**

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} & \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37}, t)} \end{array} \right] G_{16} \quad *66$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[ \begin{array}{ccc} (a'_{17})^{(2)} \boxed{+(a''_{17})^{(2)}(T_{17}, t)} & \boxed{+(a''_{14})^{(1,1)}(T_{14}, t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)} \end{array} \right] G_{17} \quad *67$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} & \boxed{+(a''_{15})^{(1,1)}(T_{14}, t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)} \end{array} \right] G_{18} \quad *68$$

Where  $+(a''_{16})^{(2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2)}(T_{17}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1)}(T_{14}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3 \*69

70

$+(a''_{36})^{(7,7)}(T_{37}, t)$ ,  $+(a''_{37})^{(7,7)}(T_{37}, t)$ ,  $+(a''_{38})^{(7,7)}(T_{37}, t)$  ARE SEVENTH DETRITION COEFFICIENTS \*71

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[ \begin{array}{ccc} (b'_{16})^{(2)} \boxed{-(b''_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{16} \quad *72$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[ \begin{array}{l} (b'_{17})^{(2)} \boxed{-(b''_{17})^{(2)}(G_{19}, t)} \quad \boxed{-(b''_{14})^{(1,1)}(G, t)} \quad \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{17} *73$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[ \begin{array}{l} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} \quad \boxed{-(b''_{15})^{(1,1)}(G, t)} \quad \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \end{array} \right] T_{18} *74$$

where  $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$  ,  $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$  ,  $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{13})^{(1,1)}(G, t)}$  ,  $\boxed{-(b''_{14})^{(1,1)}(G, t)}$  ,  $\boxed{-(b''_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$  ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$  ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$  ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$  ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1,2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$  ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$  ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1,2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$  ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$  ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detritions coefficients for category 1,2 and 3  
 $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$   $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$   $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$  are seventh detrition coefficients

**THIRD MODULE CONCATENATION: \*75**

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[ \begin{array}{l} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} \quad \boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \quad \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{20} *76$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[ \begin{array}{l} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} \quad \boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \quad \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{21} *77$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[ \begin{array}{l} (a'_{22})^{(3)} \boxed{+(a''_{22})^{(3)}(T_{21}, t)} \quad \boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \quad \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)} \end{array} \right] G_{22} *78$$

$\boxed{+(a''_{20})^{(3)}(T_{21}, t)}$  ,  $\boxed{+(a''_{21})^{(3)}(T_{21}, t)}$  ,  $\boxed{+(a''_{22})^{(3)}(T_{21}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)}$  ,  $\boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)}$  ,  $\boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)}$  are second augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$  ,  $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$  ,  $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$  are third augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$  ,  $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$  ,  $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$  ,  $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$  ,  $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$  ,  $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$  ,  $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3 \*79

80

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$   $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$   $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficient \*81



$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[ \begin{array}{l} (b'_{20})^{(3)} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} \boxed{-(b''_{36})^{(7,7,7)}(G_{19}, t)} \boxed{-(b''_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{20} \quad *82$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[ \begin{array}{l} (b'_{21})^{(3)} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{21} \quad *83$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{l} (b'_{22})^{(3)} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} \end{array} \right] T_{22} \quad *84$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$  are first detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$  are second detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$  are third detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detritions coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detritions coefficients for category 1, 2 and 3 \*85  
 $\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detritions coefficients

**FOURTH MODULE CONCATENATION: \*86**

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{24})^{(4)} \boxed{+(a''_{24})^{(4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{24} \quad *87$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{l} (a'_{25})^{(4)} \boxed{+(a''_{25})^{(4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{25} \quad *88$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{26})^{(4)} \boxed{+(a''_{26})^{(4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{26} \quad *89$$

Where  $\boxed{(a''_{24})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{25})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{26})^{(4)}(T_{25}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5)}(T_{29}, t)}$  are second augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3  
 $\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3  
 $\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, and 3  
 $\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$  ARE SEVENTH augmentation coefficients \*90

91

\*92

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{l} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) \quad - (b''_{28})^{(5,5)}(G_{31}, t) \quad - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) \quad - (b''_{16})^{(2,2,2,2)}(G_{19}, t) \quad - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,,)}(G_{39}, t) \end{array} \right] T_{24} \quad *93$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{l} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) \quad - (b''_{29})^{(5,5)}(G_{31}, t) \quad - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) \quad - (b''_{17})^{(2,2,2,2)}(G_{19}, t) \quad - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,,)}(G_{39}, t) \end{array} \right] T_{25} \quad *94$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{l} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) \quad - (b''_{30})^{(5,5)}(G_{31}, t) \quad - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) \quad - (b''_{18})^{(2,2,2,2)}(G_{19}, t) \quad - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,,)}(G_{39}, t) \end{array} \right] T_{26} \quad *95$$

Where  $-(b''_{24})^{(4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4)}(G_{27}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{32})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6)}(G_{35}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2 and 3  
 $-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2 and 3  
 $-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2 and 3  
 $-(b''_{36})^{(7,7,7,7,,)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,,)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,,)}(G_{39}, t)$  ARE SEVENTH DETRITION COEFFICIENTS\*96

\*97

**FIFTH MODULE CONCATENATION:\*98**

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \quad + (a''_{24})^{(4,4)}(T_{25}, t) \quad + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) \quad + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) \quad + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,,7,,7,7)}(T_{37}, t) \end{array} \right] G_{28} \quad *99$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{l} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) \quad + (a''_{25})^{(4,4)}(T_{25}, t) \quad + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) \quad + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) \quad + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,,7,,7,7)}(T_{37}, t) \end{array} \right] G_{29} \quad *100$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) \quad + (a''_{26})^{(4,4)}(T_{25}, t) \quad + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) \quad + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) \quad + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,,7,,7,7)}(T_{37}, t) \end{array} \right] G_{30} \quad *101$$

Where  $+(a''_{28})^{(5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5)}(T_{29}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 And  $+(a''_{24})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4)}(T_{25}, t)$  are second augmentation coefficient for category 1, 2 and 3  
 $+(a''_{32})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3  
 $+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$  are fourth augmentation coefficients for category 1, 2, and 3  
 $+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$  are fifth augmentation coefficients for category 1, 2, and 3  
 $+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$  are sixth augmentation coefficients for category 1, 2, 3 \*102

\*103

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{l} (b'_{28})^{(5)} \boxed{-(b''_{28})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{24})^{(4,4)}(G_{23}, t)} \quad \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{28} *104$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{l} (b'_{29})^{(5)} \boxed{-(b''_{29})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} \quad \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{29} *105$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{l} (b'_{30})^{(5)} \boxed{-(b''_{30})^{(5)}(G_{31}, t)} \quad \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} \quad \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{38}, t)} \end{array} \right] T_{30} *106$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1,2, and 3\*107

**SIXTH MODULE CONCATENATION\*108**

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[ \begin{array}{l} (a'_{32})^{(6)} \boxed{+(a''_{32})^{(6)}(T_{33}, t)} \quad \boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{32} *109$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{l} (a'_{33})^{(6)} \boxed{+(a''_{33})^{(6)}(T_{33}, t)} \quad \boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{33} *110$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{l} (a'_{34})^{(6)} \boxed{+(a''_{34})^{(6)}(T_{33}, t)} \quad \boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} \quad \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)} \end{array} \right] G_{34} *111$$

$\boxed{+(a''_{32})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6)}(T_{33}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$  are second augmentation coefficients for category 1,2 and 3

$\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)}$  ARE SVENTH AUGMENTATION COEFFICIENTS\*112

\*113

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{l} (b'_{32})^{(6)} \boxed{-(b''_{32})^{(6)}(G_{35}, t)} \quad \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)} \end{array} \right] T_{32} *114$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{l} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) - (b''_{29})^{(5,5,5)}(G_{31}, t) - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) \end{array} \right] T_{33} \quad *115$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{l} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) - (b''_{30})^{(5,5,5)}(G_{31}, t) - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) \end{array} \right] T_{34} \quad *116$$

$-(b''_{32})^{(6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6)}(G_{35}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4)}(G_{27}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2, and 3  
 $-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2, and 3  
 $-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2, and 3  
 $-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$  ARE SEVENTH DETRITION COEFFICIENTS\*117

\*118  
**SEVENTH MODULE CONCATENATION:**\*119

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[ (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right] + (a''_{16})^{(7)}(T_{17}, t) + (a''_{20})^{(7)}(T_{21}, t) + (a''_{24})^{(7)}(T_{23}, t)G_{36} + a_{28}''7T_{29,t} + a_{32}''7T_{33,t} + a_{13}''7T_{14,t}G_{36} \quad *120$$

121

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[ (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) \right] + (a''_{14})^{(7)}(T_{14}, t) + (a''_{21})^{(7)}(T_{21}, t) + (a''_{17})^{(7)}(T_{17}, t) + a_{25}''7T_{25,t} + a_{33}''7T_{33,t} + a_{29}''7T_{29,t} \quad G_{37}$$

\*122

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[ (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) \right] + (a''_{15})^{(7)}(T_{14}, t) + (a''_{22})^{(7)}(T_{21}, t) + (a''_{18})^{(7)}(T_{17}, t) + a_{26}''7T_{25,t} + a_{34}''7T_{33,t} + a_{30}''7T_{29,t} \quad G_{38}$$

\*123

124

125

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[ (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) - (b''_{16})^{(7)}(G_{19}, t) - (b''_{13})^{(7)}(G_{14}, t) - (b''_{20})^{(7)}(G_{231}, t) - (b''_{25})^{(7)}(G_{27}, t) - (b''_{29})^{(7)}(G_{31}, t) - (b''_{33})^{(7)}(G_{35}, t) \right] T_{36}$$

\*126

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[ (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) - (b''_{17})^{(7)}(G_{19}, t) - (b''_{19})^{(7)}(G_{14}, t) - (b''_{21})^{(7)}(G_{231}, t) - (b''_{25})^{(7)}(G_{27}, t) - (b''_{29})^{(7)}(G_{31}, t) - (b''_{33})^{(7)}(G_{35}, t) \right] T_{37}$$

\*127

Where we suppose

(A)  $(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0,$

$i, j = 13, 14, 15$

(B) The functions  $(a_i'')^{(1)}, (b_i'')^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

(C)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14}' - T_{14}| e^{-(M_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} |G' - G| e^{-(M_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T_{14}', t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T_{14}', t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

(D)  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

(E) There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together with  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\begin{aligned} \frac{dT_{38}}{dt} = & (b_{38})^{(7)} T_{37} - \left[ (b_{38})^{(7)} - \frac{(b_{38})^{(7)}((G_{39}), t)}{b26''7G27,t} - \frac{(b_{18})^{(7)}((G_{19}), t)}{-b30''7G31,t} - \frac{(b_{20})^{(7)}((G_{14}), t)}{-b34''7G35,t} \right] - \end{aligned}$$

128  
129  
130  
131  
132

$$+ (a_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \quad 134$$

$$(1)(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18 \quad 135$$

(F) (2) The functions  $(a_i'')^{(2)}, (b_i'')^{(2)}$  are positive continuous increasing and bounded. 136

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ : 137

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 138$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 139$$

(G) (3)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)}$  140

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \tag{141}$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  : 142

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition: 143

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(M_{16})^{(2)}t} \tag{144}$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(M_{16})^{(2)}t} \tag{145}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(2)}(T_{17}', t)$  and  $(a_i'')^{(2)}(T_{17}, t)$ .  $(T_{17}', t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a_i'')^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a_i'')^{(2)}(T_{17}, t)$ , the SECOND augmentation coefficient would be absolutely continuous. 146

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  : 147

(H) (4)  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants 148

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$  : 149

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \tag{150}$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \tag{151}$$

Where we suppose 152

(I) (5)  $(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, i, j = 20, 21, 22$  153

The functions  $(a_i'')^{(3)}, (b_i'')^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \tag{154}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)} \tag{155}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  : 156

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition: 157

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21}' - T_{21}| e^{-(M_{20})^{(3)}t} \tag{158}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G_{23}' - G_{23}| e^{-(M_{20})^{(3)}t} \tag{159}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(3)}(T_{21}', t)$  and  $(a_i'')^{(3)}(T_{21}, t)$ .  $(T_{21}', t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a_i'')^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a_i'')^{(3)}(T_{21}, t)$ , the THIRD augmentation coefficient, would be absolutely continuous. 160

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  : 161

(J) (6)  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , 162

satisfy the inequalities 163

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \tag{164}$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \tag{165}$$

Where we suppose 166

(K) (7)  $(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, i, j = 24, 25, 26$  167

(L) (7) The functions  $(a_i'')^{(4)}, (b_i'')^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}(G_{27}, t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$(M) \quad (8) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$  :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

171

They satisfy Lipschitz condition:

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| \leq (\hat{k}_{24})^{(4)} |(G_{27})' - (G_{27})| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(4)}(T_{25}', t)$  and  $(a_i'')^{(4)}(T_{25}, t)$ .  $(T_{25}', t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a_i'')^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 4$  then the function  $(a_i'')^{(4)}(T_{25}, t)$ , the **FOURTH augmentation coefficient WOULD** be absolutely continuous.

172

173

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$  :

174

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} + \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$  :

175

(P) (9) There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

176

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

177

(R) (10) The functions  $(a_i'')^{(5)}, (b_i'')^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$  :

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

178

$$(S) \quad (11) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$  :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition:

179

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| \leq (\hat{k}_{28})^{(5)} |(G_{31})' - (G_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(5)}(T_{29}', t)$  and  $(a_i'')^{(5)}(T_{29}, t)$ .  $(T_{29}', t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a_i'')^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 5$  then the function  $(a_i'')^{(5)}(T_{29}, t)$ , the **FIFTH augmentation coefficient** attributable would be absolutely continuous.

180

181

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  :

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} + \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$  :

182

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

183

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

184

(12) The functions  $(a_i'')^{(6)}, (b_i'')^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

185

$$(13) \lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$ :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

186

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33}' - T_{33}| e^{-(M_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} \|(G_{35})' - (G_{35})\| e^{-(M_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(6)}(T_{33}', t)$  and  $(a_i'')^{(6)}(T_{33}, t)$ .  $(T_{33}', t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a_i'')^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 6$  then the function  $(a_i'')^{(6)}(T_{33}, t)$ , the **SIXTH augmentation coefficient** would be absolutely continuous.

187

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ :

188

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$ :

189

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

190

$$(V) \quad (a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0,$$

$$i, j = 36, 37, 38$$

191

(W) The functions  $(a_i'')^{(7)}, (b_i'')^{(7)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

192

$$(X) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

**Definition of**  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$ :

Where  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$  are positive constants and  $i = 36, 37, 38$

They satisfy Lipschitz condition:

193

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(M_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), (T_{39}))| < (\hat{k}_{36})^{(7)} \|(G_{39})' - (G_{39})\| e^{-(M_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(7)}(T_{37}', t)$  and  $(a_i'')^{(7)}(T_{37}, t)$ .  $(T_{37}', t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$ . It is to be noted that  $(a_i'')^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{36})^{(7)} = 7$  then the function  $(a_i'')^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

194

**Definition of**  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ :

195



(Y)  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  :

196

(Z) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants  $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

197

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(M_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(M_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

198

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(M_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(M_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

199

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 200

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 201$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(M_{13})^{(1)}t} \quad 202$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(M_{13})^{(1)}t} \quad 203$$

By 204

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t [(a_{13})^{(1)} G_{14}(s_{(13)}) - ((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})] G_{13}(s_{(13)})] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t [(a_{14})^{(1)} G_{13}(s_{(13)}) - ((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}))] G_{14}(s_{(13)})] ds_{(13)} \quad 205$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t [(a_{15})^{(1)} G_{14}(s_{(13)}) - ((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}))] G_{15}(s_{(13)})] ds_{(13)} \quad 206$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t [(b_{13})^{(1)} T_{14}(s_{(13)}) - ((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{13}(s_{(13)})] ds_{(13)} \quad 207$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t [(b_{14})^{(1)} T_{13}(s_{(13)}) - ((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{14}(s_{(13)})] ds_{(13)} \quad 208$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t [(b_{15})^{(1)} T_{14}(s_{(13)}) - ((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}))] T_{15}(s_{(13)})] ds_{(13)} \quad 209$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

210

if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t}$$

By

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36}(s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + a''_{37}(s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + a''_{38}(s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 211

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)}, \quad 212$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(M_{16})^{(2)}t} \quad 213$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(M_{16})^{(2)}t} \quad 214$$

By 215

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16}(s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + a''_{17}(s_{(16)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)} \quad 216$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + a''_{18}(s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)} \quad 217$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)} \quad 218$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)} \quad 219$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)} \quad 220$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

221

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)}, \quad 222$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(M_{20})^{(3)}t} \quad 223$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(M_{20})^{(3)}t} \quad 224$$

By 225

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20}(s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)} \quad 226$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)} \quad 227$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)} \quad 228$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)} \quad 229$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)} \quad 230$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)} \quad 231$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 231

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)}, \quad 232$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(M_{24})^{(4)}t} \quad 233$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(M_{24})^{(4)}t} \quad 234$$

By 235

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24}(s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)} \quad 236$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)} \quad 237$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)} \quad 238$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)} \quad 239$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)} \quad 240$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)} \quad 241$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 241

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)}, \quad 242$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(M_{28})^{(5)}t} \quad 243$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(M_{28})^{(5)}t} \quad 244$$

By 245

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28}(s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)} \quad 246$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)} \quad 247$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)} \quad 248$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)} \quad 249$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)} \quad 250$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)} \quad 251$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

252

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)}, \tag{253}$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(M_{32})^{(6)}t} \tag{254}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(M_{32})^{(6)}t} \tag{255}$$

By 256

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32}(s_{(32)}) \right) T_{33}(s_{(32)}) \right] G_{32}(s_{(32)}) ds_{(32)} \tag{257}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + a''_{33}(s_{(32)}) \right) T_{33}(s_{(32)}) \right] G_{33}(s_{(32)}) ds_{(32)} \tag{258}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + a''_{34}(s_{(32)}) \right) T_{33}(s_{(32)}) \right] G_{34}(s_{(32)}) ds_{(32)} \tag{259}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)}(G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)} \tag{260}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)}(G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)} \tag{261}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)}(G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)} \tag{262}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

: if the conditions IN THE FOREGOING are fulfilled, there exists a solution satisfying the conditions 262

**Definition of  $G_i(0), T_i(0)$  :**

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)}, \tag{263}$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t} \tag{264}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t} \tag{265}$$

By 266

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36}(s_{(36)}) \right) T_{37}(s_{(36)}) \right] G_{36}(s_{(36)}) ds_{(36)} \tag{267}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + a''_{37}(s_{(36)}) \right) T_{37}(s_{(36)}) \right] G_{37}(s_{(36)}) ds_{(36)} \tag{268}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + a''_{38}(s_{(36)}) \right) T_{37}(s_{(36)}) \right] G_{38}(s_{(36)}) ds_{(36)} \tag{269}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)} \tag{270}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)} \tag{271}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)}(G(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)} \tag{272}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$  272

(a) The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that 273

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(M_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(P_{24})^{(4)}}{(M_{24})^{(4)}}(e^{(M_{24})^{(4)}t} - 1)$$

From which it follows that

274

$$(G_{24}(t) - G_{24}^0)e^{-(M_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(M_{24})^{(4)}} \left[ ((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{((\hat{P}_{24})^{(4)} + G_{25}^0)}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(b) The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that 275

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(M_{28})^{(5)}s(28)} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(P_{28})^{(5)}}{(M_{28})^{(5)}}(e^{(M_{28})^{(5)}t} - 1)$$

From which it follows that

276

$$(G_{28}(t) - G_{28}^0)e^{-(M_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(M_{28})^{(5)}} \left[ ((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{((\hat{P}_{28})^{(5)} + G_{29}^0)}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

(c) The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that 277

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(M_{32})^{(6)}s(32)} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(P_{32})^{(6)}}{(M_{32})^{(6)}}(e^{(M_{32})^{(6)}t} - 1)$$

From which it follows that

278

$$(G_{32}(t) - G_{32}^0)e^{-(M_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(M_{32})^{(6)}} \left[ ((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{((\hat{P}_{32})^{(6)} + G_{33}^0)}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(d) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying 37,35,36 into itself .Indeed it is obvious that 279

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} \left( G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}s(36)} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(P_{36})^{(7)}}{(M_{36})^{(7)}}(e^{(M_{36})^{(7)}t} - 1)$$

From which it follows that

280

$$(G_{36}(t) - G_{36}^0)e^{-(M_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(M_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{((\hat{P}_{36})^{(7)} + G_{37}^0)}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(M_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(M_{13})^{(1)}} < 1$  and to choose 281

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have 282

$$\frac{(a_i)^{(1)}}{(M_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0)e^{-\frac{((\hat{P}_{13})^{(1)} + G_j^0)}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)} \quad 283$$

$$\frac{(b_i)^{(1)}}{(M_{13})^{(1)}} \left[ ((\hat{Q}_{13})^{(1)} + T_j^0)e^{-\frac{((\hat{Q}_{13})^{(1)} + T_j^0)}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \quad 284$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying GLOBAL EQUATIONS into itself 285

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 286

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(M_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(M_{13})^{(1)}t} \}$$

Indeed if we denote 287

**Definition of  $\tilde{G}, \tilde{T}$  :**

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + G_{13}^{(2)} |(a'_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} \leq \frac{1}{(\bar{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}) d((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})) \tag{288}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  and  $(\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ . 289

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$ ,  $i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  290

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0 \tag{291}$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\bar{M}_{13})^{(1)})_1$ , and  $((\bar{M}_{13})^{(1)})_3$  : 292

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$G_{13} < (\bar{M}_{13})^{(1)}$  it follows  $\frac{dG_{14}}{dt} \leq ((\bar{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$  and by integrating

$$G_{14} \leq ((\bar{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\bar{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\bar{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\bar{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$ ,  $G_{15}$  and  $G_{13}$ ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 293

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$  then  $T_{14} \rightarrow \infty$ . 294

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to 295

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded. The}$$

same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} ((b_{15}'')^{(1)}(G(t), t)) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions 296

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$  and to choose 297

$(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \tag{298}$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \tag{299}$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 300

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric 301

$$d(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

$$\underline{\text{Definition of}} \widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$$

It results

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(2)}$  and  $(b''_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(2)} - (a''_i)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)}]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $((\bar{M}_{16})^{(2)})_1, ((\bar{M}_{16})^{(2)})_2$  and  $((\bar{M}_{16})^{(2)})_3$  :

**Remark 3:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$$\begin{aligned} G_{16} < ((\bar{M}_{16})^{(2)}) &\text{ it follows } \frac{dG_{17}}{dt} \leq ((\bar{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17} \text{ and by integrating} \\ G_{17} &\leq ((\bar{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\bar{M}_{16})^{(2)})_1 / (a'_{17})^{(2)} \end{aligned}$$

In the same way, one can obtain

$$G_{18} \leq ((\bar{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\bar{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively.

**Remark 4:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below.

**Remark 5:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(2)} ((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ .

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b''_i)^{(2)} ((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)} (m)^{(2)} - \varepsilon_2 T_{17}$  which leads to

$$T_{17} \geq \left( \frac{(a_{17})^{(2)} (m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left( \frac{(a_{17})^{(2)} (m)^{(2)}}{2} \right), t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded. The}$$

same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b''_{18})^{(2)} ((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(M_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(M_{20})^{(3)}} < 1$  and to choose

$(\widehat{P}_{20})^{(3)}$  and  $(\widehat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(M_{20})^{(3)}} \left[ (\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)}$$

$$\frac{(b_i)^{(3)}}{(M_{20})^{(3)}} \left[ ((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)}$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric

320

$$d\left(\left((G_{23})^{(1)}, (T_{23})^{(1)}\right), \left((G_{23})^{(2)}, (T_{23})^{(2)}\right)\right) = \sup_{t \in \mathbb{R}_+} \left\{ \max_i |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_i |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote

321

**Definition of**  $\widetilde{G}_{23}, \widetilde{T}_{23} : ((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

322

$$|\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} +$$

$$\int_0^t \left\{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \right.$$

323

$$(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + G_{20}^{(2)} |(a'_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a'_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

324

$$|G^{(1)} - G^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} \leq$$

$$\frac{1}{(\bar{M}_{20})^{(3)}} \left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} + (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} \right) d\left(\left((G_{23})^{(1)}, (T_{23})^{(1)}\right); (G_{23})^{(2)}, (T_{23})^{(2)}\right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

325

**Remark 1:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  and  $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$ ,  $i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

326

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \left\{ (a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0$$

327

**Definition of**  $((\bar{M}_{20})^{(3)})_1, ((\bar{M}_{20})^{(3)})_2$  and  $((\bar{M}_{20})^{(3)})_3$  :

**Remark 3:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$ . indeed if

$$G_{20} < (\bar{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\bar{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\bar{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\bar{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\bar{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\bar{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$ ,  $G_{22}$  and  $G_{20}$ ,  $G_{21}$  respectively.

328

**Remark 4:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below.

329

**Remark 5:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$  then  $T_{21} \rightarrow \infty$ .

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

330

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21})^{(3)} - (b''_i)^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

331

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded. The}$$

same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b''_{22})^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

332

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$  and to choose

333

$(\bar{P}_{24})^{(4)}$  and  $(\bar{Q}_{24})^{(4)}$  large to have



$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ (\hat{P}_{24})^{(4)} + ((\hat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{24})^{(4)} \quad 334$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ ((\hat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)} \quad 335$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying IN to itself 336

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 337

$$d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\overline{G_{27}}), (\overline{T_{27}})$  :  $((\overline{G_{27}}), (\overline{T_{27}})) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s(24)} e^{(\bar{M}_{24})^{(4)}s(24)} ds(24) + \\ &\int_0^t \{ (a_{24}')^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s(24)} e^{-(\bar{M}_{24})^{(4)}s(24)} + \\ &(a_{24}'')^{(4)} (T_{25}^{(1)}, s(24)) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s(24)} e^{(\bar{M}_{24})^{(4)}s(24)} + \\ &G_{24}^{(2)} |(a_{24}'')^{(4)} (T_{25}^{(1)}, s(24)) - (a_{24}'')^{(4)} (T_{25}^{(2)}, s(24))| e^{-(\bar{M}_{24})^{(4)}s(24)} e^{(\bar{M}_{24})^{(4)}s(24)} \} ds(24) \end{aligned}$$

Where  $s(24)$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows 338

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{1}{(\bar{M}_{24})^{(4)}} \left( (a_{24})^{(4)} + (a_{24}')^{(4)} + (\bar{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{K}_{24})^{(4)} \right) d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}); (G_{27})^{(2)}, (T_{27})^{(2)} \right) \quad 339$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a_{24}'')^{(4)}$  and  $(b_{24}'')^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  and  $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 340

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  341

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a_i')^{(4)} - (a_i'')^{(4)} (T_{25}(s(24)), s(24)) \} ds(24)} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\bar{M}_{24})^{(4)})_1, ((\bar{M}_{24})^{(4)})_2$  and  $((\bar{M}_{24})^{(4)})_3$  : 342

**Remark 3:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$  . indeed if

$G_{24} < (\bar{M}_{24})^{(4)}$  it follows  $\frac{dG_{25}}{dt} \leq ((\bar{M}_{24})^{(4)})_1 - (a_{25}')^{(4)} G_{25}$  and by integrating

$$G_{25} \leq ((\bar{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\bar{M}_{24})^{(4)})_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq ((\bar{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\bar{M}_{24})^{(4)})_2 / (a_{26}')^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$  ,  $G_{26}$  and  $G_{24}$  ,  $G_{25}$  respectively.

**Remark 4:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below. 343

**Remark 5:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(4)} ((G_{27})(t), t)) = (b_{25}')^{(4)}$  then  $T_{25} \rightarrow \infty$ . 344

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)} ((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)} (m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 345

$$T_{25} \geq \left( \frac{(a_{25})^{(4)} (m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$T_{25} \geq \left( \frac{(a_{25})^{(4)} (m)^{(4)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_4}$  By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded. The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)} ((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities

hold also for  $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

346

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$  and to choose  $(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

347

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ (\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\widehat{P}_{28})^{(5)}$$

348

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ ((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)}$$

349

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

350

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric

351

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t} \right\}$$

352

Indeed if we denote

$$\text{Definition of } (\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$$

It results

$$\begin{aligned} |\widehat{G}_{28}^{(1)} - \widehat{G}_{28}^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} &\left( (a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)} \right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right); (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned}$$

353

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (35,35,36) the result follows

**Remark 1:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ .

354

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}, i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

355

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(5)} - (a''_i)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

$$\text{Definition of } ((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2 \text{ and } ((\widehat{M}_{28})^{(5)})_3 :$$

356

**Remark 3:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)} ((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}, G_{30}$  and  $G_{28}, G_{29}$  respectively.

**Remark 4:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below.

357

**Remark 5:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$  then  $T_{29} \rightarrow \infty$ .

358

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b''_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to

360

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$$

If we take  $t$  such that  $e^{-\varepsilon_5 t} = \frac{1}{2}$  it results

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5}$$

By taking now  $\varepsilon_5$  sufficiently small one sees that  $T_{29}$  is unbounded. The

same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b_{30}'')^{(5)}((G_{31})(t), t) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

361

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(M_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(M_{32})^{(6)}} < 1$  and to choose

362

$(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(M_{32})^{(6)}} \left[ (\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)}$$

363

$$\frac{(b_i)^{(6)}}{(M_{32})^{(6)}} \left[ ((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)}$$

364

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself

365

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric

366

$$d\left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(M_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(M_{32})^{(6)}t} \right\}$$

Indeed if we denote

$$\underline{\text{Definition of}} \left( \widehat{G}_{35}, \widehat{T}_{35} \right) : \left( \widehat{G}_{35}, \widehat{T}_{35} \right) = \mathcal{A}^{(6)}\left( (G_{35}), (T_{35}) \right)$$

It results

$$\begin{aligned} |\widehat{G}_{32}^{(1)} - \widehat{G}_{32}^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(M_{32})^{(6)}s} e^{(M_{32})^{(6)}s} ds + \\ &\int_0^t \{ (a_{32}')^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(M_{32})^{(6)}s} e^{-(M_{32})^{(6)}s} + \\ &(a_{32}'')^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(M_{32})^{(6)}s} e^{(M_{32})^{(6)}s} + \\ &G_{32}^{(2)} | (a_{32}'')^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a_{32}'')^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(M_{32})^{(6)}s} e^{(M_{32})^{(6)}s} \} ds \end{aligned}$$

367

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$(1) \quad (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \\ i, j = 13, 14, 15$$

(2) The functions  $(a_i'')^{(1)}, (b_i'')^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$(3) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants  
 and  $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T_{14}'| e^{-(M_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, T)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(M_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the **first augmentation coefficient** attributable to terrestrial organisms, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

(AA)  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

(BB) There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together with  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Analogous inequalities hold also for  $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$

368

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 7$  and to choose  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ (\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{36})^{(7)}$$

369

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ ((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)}$$

370

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 37,35,36 into itself

371

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric

372

$$d\left((G_{39})^{(1)}, (T_{39})^{(1)}, (G_{39})^{(2)}, (T_{39})^{(2)}\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{39}), (\widehat{T}_{39})$  :

$$((\widehat{G}_{39}), (\widehat{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_{36}^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s} e^{(\hat{M}_{36})^{(7)}s} ds + \\ &\int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s} e^{-(\hat{M}_{36})^{(7)}s} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\hat{M}_{36})^{(7)}s} e^{(\hat{M}_{36})^{(7)}s} + \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

373

$$\begin{aligned} & |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widehat{M}_{36})^{(7)}t} \leq \\ & \frac{1}{(\widehat{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a_{36}')^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}) d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (37,35,36) the result follows 374

**Remark 1:** The fact that we supposed  $(a_{36}'')^{(7)}$  and  $(b_{36}'')^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 375

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

376

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 79 to 36 it results

$$\begin{aligned} G_i(t) & \geq G_i^0 e^{-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0 \\ T_i(t) & \geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \quad \text{for } t > 0 \end{aligned}$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  : 377

**Remark 3:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$  it follows  $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$ ,  $G_{38}$  and  $G_{36}$ ,  $G_{37}$  respectively.

**Remark 7:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 378

**Remark 5:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$  then  $T_{37} \rightarrow \infty$ . 379

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 380

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded. The same property holds for } T_{38} \text{ if } \lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 72

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying GLOBAL EQUATIONS AND ITS CONCOMITANT CONDITIONALITIES into itself 381

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 382

$$d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(M_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(M_{36})^{(7)}t} \}$$

Indeed if we denote

**Definition of**  $(\widetilde{G_{39}}), (\widetilde{T_{39}})$  :

$$((\widetilde{G_{39}}), (\widetilde{T_{39}})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$$

It results

$$\begin{aligned} |\widetilde{G}_{36}^{(1)} - \widetilde{G}_{36}^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(M_{36})^{(7)}s_{(36)}} e^{(M_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a_{36}')^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(M_{36})^{(7)}s_{(36)}} e^{-(M_{36})^{(7)}s_{(36)}} + \\ &(a_{36}'')^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(M_{36})^{(7)}s_{(36)}} e^{(M_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a_{36}'')^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a_{36}'')^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(M_{36})^{(7)}s_{(36)}} e^{(M_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows 384

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(M_{36})^{(7)}t} &\leq \\ \frac{1}{(M_{36})^{(7)}} &((a_{36})^{(7)} + (a_{36}')^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a_{36}'')^{(7)}$  and  $(b_{36}'')^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(M_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 385

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}, i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  386

From CONCATENATED GLOBAL EQUATIONS it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{-\int_0^t \{ (a_i')^{(7)} - (a_i'')^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b_i')^{(7)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  : 387

**Remark 3:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$$G_{36} < (\widehat{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a_{37}')^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)} ((\widehat{M}_{36})^{(7)})_1 / (a_{37}')^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)} ((\widehat{M}_{36})^{(7)})_2 / (a_{38}')^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$ ,  $G_{38}$  and  $G_{36}$ ,  $G_{37}$  respectively.

**Remark 7:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 388

**Remark 5:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$  then  $T_{37} \rightarrow \infty$ . 389

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 390

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_7}$  By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is unbounded. The

same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \quad 391$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 392$$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  : 393

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots 394

(a) of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  395

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and 396

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  : 397

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the 398

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  399

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  400

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :- 401

(b) If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by 402

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 403$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 404$$

and  $(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 405$$

and analogously 406

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and  $(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 407$$

Then the solution satisfies the inequalities 408

$$G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$  is defined 409

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 410$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{(S_1)^{(2)} - (p_{16})^{(2)} t} - e^{-(S_2)^{(2)} t} \right] + G_{18}^0 e^{-(S_2)^{(2)} t} \right) \leq G_{18}(t) \leq \quad 411$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a_{18})^{(2)})} \left[ e^{(S_1)^{(2)} t} - e^{-(a_{18})^{(2)} t} \right] + G_{18}^0 e^{-(a_{18})^{(2)} t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq T_{16}^0 e^{(R_1)^{(2)} + (r_{16})^{(2)} t} \quad 412$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} + (r_{16})^{(2)} t} \quad 413$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b_{18})^{(2)})} \left[ e^{(R_1)^{(2)} t} - e^{-(b_{18})^{(2)} t} \right] + T_{18}^0 e^{-(b_{18})^{(2)} t} \leq T_{18}(t) \leq \quad 414$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{(R_1)^{(2)} + (r_{16})^{(2)} t} - e^{-(R_2)^{(2)} t} \right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$  :- 415

Where  $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$  416

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 417$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

418  
419

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

- (a)  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying  
 $-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$   
 $-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

- (b) By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

- (c) If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by  
 $(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$ , **if**  $(v_0)^{(3)} < (v_1)^{(3)}$   
 $(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$ , **if**  $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$ ,

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$ , **if**  $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously

$$(u_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$$
, **if**  $(u_0)^{(3)} < (u_1)^{(3)}$

$$(u_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$$
, **if**  $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$ , and  $(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$

$(u_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$ , **if**  $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$$

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b_{22})^{(3)}t} \leq T_{22}(t) \leq$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[ e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$  :-

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b_{22})^{(3)} - (r_{22})^{(3)}$$

430  
431  
432

If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

- (d)  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$



$$-(\tau_2)^{(4)} \leq -(b_{24}')^{(4)} + (b_{25}')^{(4)} - (b_{24}'')^{(4)}((G_{27}), t) - (b_{25}'')^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 433

(e) By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 434

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$
 436

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

(f) If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously 437

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)}$  where  $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$  are defined respectively

Then the solution satisfies the inequalities 439

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined 440

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$
 441

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq (a_{26})^{(4)} G_{24}^0 (m_2)^{(4)} (S_1)^{(4)} - (a_{26}')^{(4)} e^{(S_1)^{(4)}t} + G_{26}^0 e^{-(a_{26}')^{(4)}t}$$
 442

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$
 443

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$
 444

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b_{26}')^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b_{26}')^{(4)}t} \right] + T_{26}^0 e^{-(b_{26}')^{(4)}t} \leq T_{26}(t) \leq$$
 445

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})} \left[ e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :- 452

Where  $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)} \tag{453}$$

**Behavior of the solutions** 454

If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

(g)  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  : 455

(h) By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  : 456

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the

roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

(i) If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{a_{28}^0}{a_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 457

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)}$  where  $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$  are defined respectively

Then the solution satisfies the inequalities 458

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 459$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \right) \leq G_{30}(t) \leq \quad 460$$

$$(a_{30})^{(5)} G_{28}^0 (m_2)^{(5)} (S_1)^{(5)} - (a_{30})^{(5)} e^{(S_1)^{(5)}t} - e^{-(a_{30})^{(5)}t} + G_{30}^0 e^{-(a_{30})^{(5)}t} \quad 461$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 462$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 463$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 464$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t} \quad 465$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 465

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions** 466

If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

(j)  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 467

(k) By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 468

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

(l) If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

471

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)}$  where  $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$  are defined respectively

Then the solution satisfies the inequalities

472

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

473

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq (a_{34})^{(6)} G_{32}^0 (m_2)^{(6)} (S_1)^{(6)} - (a_{34}')^{(6)} e^{(S_1)^{(6)}t} + G_{34}^0 e^{-(a_{34}')^{(6)}t}$$

474

$$T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

475

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

476

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b_{34}')^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b_{34}')^{(6)}t} \right] + T_{34}^0 e^{-(b_{34}')^{(6)}t} \leq T_{34}(t) \leq$$

477

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$ :-

478

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a_{32}')^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b_{32}')^{(6)}$$

$$(R_2)^{(6)} = (b_{34}')^{(6)} - (r_{34})^{(6)}$$

479

If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

(m)  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a_{36}')^{(7)} + (a_{37}')^{(7)} - (a_{36}'')^{(7)} (T_{37}, t) + (a_{37}'')^{(7)} (T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b_{36}')^{(7)} + (b_{37}')^{(7)} - (b_{36}'')^{(7)} (G_{39}, t) - (b_{37}'')^{(7)} (G_{39}, t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  :

480

(n) By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)} (v^{(7)})^2 + (\sigma_1)^{(7)} v^{(7)} - (a_{36})^{(7)} = 0$$

481

$$\text{and } (b_{37})^{(7)} (u^{(7)})^2 + (\tau_1)^{(7)} u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  :

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

(o) If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

are defined respectively

Then the solution satisfies the inequalities

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t}$$

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \right) \leq G_{38}(t) \leq$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a_{38})^{(7)}t}$$

$$\frac{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b_{38})^{(7)}t} \leq T_{38}(t) \leq$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$  :-

$$\text{Where } (S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b_{38})^{(7)} - (r_{38})^{(7)}$$

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) -$$

**Definition of**  $v^{(7)}$  :-  $v^{(7)} = \frac{G_{36}}{G_{37}}$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

(a) For  $0 < \frac{G_{36}^0}{G_{37}^0} < (v_0)^{(7)} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad (C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

493

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}, \quad (\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

(b) If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case,

494

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (\bar{v}_1)^{(7)}$$

(c) If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \frac{G_{36}^0}{G_{37}^0}$ , we obtain

495

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :-

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{36}'' )^{(7)} = (a_{37}'' )^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b_{36}'' )^{(7)} = (b_{37}'' )^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then

$(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

We can prove the following

If  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  are independent on  $t$ , and the conditions

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined are satisfied, then the system WITH THE SATISFACTION OF THE FOLLOWING PROPERTIES HAS A SOLUTION AS DERIVED BELOW.

**Particular case :**

If  $(a''_{16})^{(2)} = (a''_{17})^{(2)}$ , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if  $(b''_{16})^{(2)} = (b''_{17})^{(2)}$ , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of  $v^{(3)}$  :-** 
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$

It follows

$$- \left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

From which one obtains

(a) For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner, we get

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of  $(\bar{v}_1)^{(3)}$  :-**

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

(b) If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case,

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

(c) If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$ , we obtain

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(3)}(t)$  :-**

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{20}''^{(3)}) = (a_{21}''^{(3)})$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b_{20}''^{(3)}) = (b_{21}''^{(3)})$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

506

: From GLOBAL EQUATIONS we obtain

507

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t) \right) - (a_{25}'')^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

508

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

$$(d) \text{ For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_0)^{(4)}]t}}, \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner, we get

509

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}, \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

$$(e) \text{ If } 0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)} \text{ we find like in the previous case,}$$

510

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_2)^{(4)}]t}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}(v_1)^{(4)} - (v_2)^{(4)}]t}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}} \leq (\bar{v}_1)^{(4)}$$

511

$$(f) \text{ If } 0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}, \text{ we obtain}$$

512

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}(\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}]t}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(4)}(t)$  :-



$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{24})^{(4)} = (a''_{25})^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$  **this also defines  $(v_0)^{(4)}$  for the special case .** 513

Analogously if  $(b''_{24})^{(4)} = (b''_{25})^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of  $(u_0)^{(4)}$ .**

From GLOBAL EQUATIONS we obtain

514  
515

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

(g) For  $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}, \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

516

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}, \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

(h) If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

517

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (C)^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}} \leq (\bar{v}_1)^{(5)}$$

(i) If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$ , we obtain

518

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (C)^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}} \leq (v_0)^{(5)}$$

519

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{28})^{(5)} = (a''_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

520

we obtain

521

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)} (T_{33}, t) \right) - (a''_{33})^{(6)} (T_{33}, t) v^{(6)} - (a_{33})^{(6)} v^{(6)}$$

**Definition of**  $v^{(6)}$  :-  $v^{(6)} = \frac{G_{32}}{G_{33}}$

It follows

$$- \left( (a_{33})^{(6)} (v^{(6)})^2 + (\sigma_2)^{(6)} v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

(j) For  $0 < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} (v_1)^{(6)} - (v_0)^{(6)}] t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)} (v_1)^{(6)} - (v_0)^{(6)}] t}}, \quad (C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner, we get

522

523

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}}, \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case, 524

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (\bar{C})^{(6)} (v_2)^{(6)} e^{[-(a_{33})^{(6)} (v_1)^{(6)} - (v_2)^{(6)}] t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} (v_1)^{(6)} - (v_2)^{(6)}] t}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}} \leq (\bar{v}_1)^{(6)}$$

(l) If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$ , we obtain 525

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)} (\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)} (\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}] t}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{32})^{(6)} = (a_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)} G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case .**

Analogously if  $(b_{32})^{(6)} = (b_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then  $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

526

**Behavior of the solutions**

527

If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

(p)  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  : 528

(q) By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

$$\text{and } (b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0 \text{ and}$$

529

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  :

530.

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

(r) If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

531

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}$ , if  $(\bar{u}_1)^{(7)} < (u_0)^{(7)}$  where  $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$  are defined by 59 and 67 respectively

Then the solution of GLOBAL EQUATIONS satisfies the inequalities

532

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t}$$

533

(

534

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\frac{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

535

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

536

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq$$

537

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :-

538

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

539

$$(R_2)^{(7)} = (b_{38})^{(7)} - (r_{38})^{(7)}$$

From CONCATENATED GLOBAL EQUATIONS we obtain

540

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :- 
$$v^{(7)} = \frac{G_{36}}{G_{37}}$$

It follows

$$-\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

(m) For  $0 < \frac{G_{36}^0}{G_{37}^0} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad (C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner, we get

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}, \quad (\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

541

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

(n) If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case,

542

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (\bar{v}_1)^{(7)}$$

(o) If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \frac{G_{36}^0}{G_{37}^0}$ , we obtain

543

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :-

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}$$

Now, using this result and replacing it in CONCATENATED GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{36})^{(7)} = (a_{37})^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case.**

Analogously if  $(b_{36})^{(7)} = (b_{37})^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then

$(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0$$

544

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0$$

545

has a unique positive solution, which is an equilibrium solution for the system

546

$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0$	547
$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0$	548
$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0$	549
$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0$	550
$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0$	551
$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0$	552
has a unique positive solution , which is an equilibrium solution for	553
$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0$	554
$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0$	555
$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0$	556
$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0$	557
$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0$	558
$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0$	559
has a unique positive solution , which is an equilibrium solution	560
$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0$	561
$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0$	563
$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0$	564
$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0$	565
$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0$	566
$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0$	567
has a unique positive solution , which is an equilibrium solution for the system	568
$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0$	569
$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0$	570
$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0$	571
$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0$	572
$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0$	573
$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0$	574
has a unique positive solution , which is an equilibrium solution for the system	575
$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0$	576
$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0$	577
$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0$	578
$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0$	579
$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0$	580

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 584$$

has a unique positive solution , which is an equilibrium solution for the system 582

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 583$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 584$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 585$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 586$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 588$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 589$$

has a unique positive solution , which is an equilibrium solution for the system 560

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Definition and uniqueness of  $T_{37}^*$**  :- 561

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(7)}(T_{37})$  being increasing, it follows that there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)} + (a_{36})^{(7)}(T_{37}^*)]} , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)} + (a_{38})^{(7)}(T_{37}^*)]}$$

(e) By the same argument, the equations ( SOLUTIONAL) admit solutions  $G_{36}, G_{37}$  if

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - [(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G^*) = 0$  562

Finally we obtain the unique solution OF THE SYSTEM

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0, T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)} + (a_{36})^{(7)}(T_{37}^*)]} , \quad G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)} + (a_{38})^{(7)}(T_{37}^*)]} \\ T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)} - (b_{36})^{(7)}((G_{39})^*)]} , \quad T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)} - (b_{38})^{(7)}((G_{39})^*)]} \quad 563$$

**Definition and uniqueness of  $T_{21}^*$**  :- 564

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a_{20})^{(3)}(T_{21}^*)]} , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $T_{25}^*$**  :- 566

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a_{24})^{(4)}(T_{25}^*)]} , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $T_{29}^*$**  :- 567

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a_{28})^{(5)}(T_{29}^*)]} , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $T_{33}^*$**  :- 568

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

(f) By the same argument, the equations 92,93 admit solutions  $G_{13}, G_{14}$  if 569

$$\varphi(G) = (b'_{13})^{(1)}(b_{14})^{(1)} - (b_{13})^{(1)}(b'_{14})^{(1)} - [(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

(g) By the same argument, the equations 92,93 admit solutions  $G_{16}, G_{17}$  if 570

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - [(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{17}^*$  such that  $\varphi((G_{19})^*) = 0$  571

(a) By the same argument, the concatenated equations admit solutions  $G_{20}, G_{21}$  if 572

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - [(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$  573

(b) By the same argument, the equations of modules admit solutions  $G_{24}, G_{25}$  if 574

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - [(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

(c) By the same argument, the equations (modules) admit solutions  $G_{28}, G_{29}$  if 575

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - [(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

(d) By the same argument, the equations (modules) admit solutions  $G_{32}, G_{33}$  if 578

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - [(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G^*) = 0$  580

Finally we obtain the unique solution of 89 to 94 582

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 583

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0, T_{17}^*$  given by  $f(T_{17}^*) = 0$  and 584

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]}$$

Obviously, these values represent an equilibrium solution 587

Finally we obtain the unique solution 588

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0, T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$



$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b_{20})^{(3)} - (b_{20}'')^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b_{22})^{(3)} - (b_{22}'')^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{25}^*$  given by  $\varphi(G_{27}) = 0$  ,  $T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a_{24})^{(4)} + (a_{24}'')^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a_{26})^{(4)} + (a_{26}'')^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b_{24})^{(4)} - (b_{24}'')^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b_{26})^{(4)} - (b_{26}'')^{(4)}((G_{27})^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$  ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a_{28})^{(5)} + (a_{28}'')^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a_{30})^{(5)} + (a_{30}'')^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b_{28})^{(5)} - (b_{28}'')^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b_{30})^{(5)} - (b_{30}'')^{(5)}((G_{31})^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$  ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a_{32})^{(6)} + (a_{32}'')^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a_{34})^{(6)} + (a_{34}'')^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b_{32})^{(6)} - (b_{32}'')^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b_{34})^{(6)} - (b_{34}'')^{(6)}((G_{35})^*)]}$$

Obviously, these values represent an equilibrium solution

**ASYMPTOTIC STABILITY ANALYSIS**

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  Belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i , T_i = T_i^* + T_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{13}}{dt} = -((a_{13}')^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14}$$

$$\frac{dG_{14}}{dt} = -((a_{14}')^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14}$$

$$\frac{dG_{15}}{dt} = -((a_{15}')^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14}$$

$$\frac{dT_{13}}{dt} = -((b_{13}')^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j$$

$$\frac{dT_{14}}{dt} = -((b_{14}')^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j$$

$$\frac{dT_{15}}{dt} = -((b_{15}')^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  Belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i , T_i = T_i^* + T_i$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} , \frac{\partial (b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij}$$

taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a_{16}')^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17}$$

$$\frac{dG_{17}}{dt} = -((a_{17}')^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17}$$

$$\frac{dG_{18}}{dt} = -((a_{18}')^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17}$$

$$\frac{dT_{16}}{dt} = -((b_{16}')^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j$$

$$\frac{dT_{17}}{dt} = -((b_{17}')^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j$$

$$\frac{dT_{18}}{dt} = -((b_{18}')^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  Belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{21}''^{(3)})}{\partial T_{21}} (T_{21}^*) = (q_{21})^{(3)}, \frac{\partial (b_i''^{(3)})}{\partial G_j} ((G_{23})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^* \mathbb{T}_{21} \tag{616}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^* \mathbb{T}_{21} \tag{617}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^* \mathbb{T}_{21} \tag{618}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^* \mathbb{G}_j \tag{619}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^* \mathbb{G}_j \tag{620}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^* \mathbb{G}_j \tag{621}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of  $\mathbb{G}_i, \mathbb{T}_i$  :-**

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}''^{(4)})}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)}, \frac{\partial (b_i''^{(4)})}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^* \mathbb{T}_{25} \tag{622}$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^* \mathbb{T}_{25} \tag{623}$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^* \mathbb{T}_{25} \tag{624}$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^* \mathbb{G}_j \tag{625}$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^* \mathbb{G}_j \tag{626}$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^* \mathbb{G}_j \tag{627}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of  $\mathbb{G}_i, \mathbb{T}_i$  :-**

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}''^{(5)})}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)}, \frac{\partial (b_i''^{(5)})}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^* \mathbb{T}_{29} \tag{628}$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^* \mathbb{T}_{29} \tag{629}$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^* \mathbb{T}_{29} \tag{630}$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^* \mathbb{G}_j \tag{631}$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^* \mathbb{G}_j \tag{632}$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^* \mathbb{G}_j \tag{633}$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of  $\mathbb{G}_i, \mathbb{T}_i$  :-**

$$G_i = G_i^* + \mathbb{G}_i, T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}''^{(6)})}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)}, \frac{\partial (b_i''^{(6)})}{\partial G_j} ((G_{35})^*) = s_{ij}$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^* \mathbb{T}_{33} \tag{634}$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^* \mathbb{T}_{33} \tag{635}$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^* \mathbb{T}_{33} \tag{636}$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^* \mathbb{G}_j \tag{637}$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^* G_j \tag{649}$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^* G_j \tag{650}$$

Obviously, these values represent an equilibrium solution of 79,20,36,22,23, 651

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 652

$$G_i = G_i^* + G_i \quad , \quad T_i = T_i^* + T_i \tag{653}$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations(SOLUTIONAL) and neglecting the terms of power 2, we obtain 654

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^* T_{37} \tag{655}$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^* T_{37} \tag{657}$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^* T_{37} \tag{658}$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^* G_j \tag{659}$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^* G_j \tag{660}$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^* G_j \tag{661}$$

$$2. \tag{662}$$

The characteristic equation of this system is

$$\begin{aligned} & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ & \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & \left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & + \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left( (a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right)\} = 0 \\ & + \\ & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ & \left[ ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ & \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ & + \left( ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ & \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ & \left( ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \\ & \left( ((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \right) \\ & + \left( ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) (q_{18})^{(2)}G_{18} \\ & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) \left( (a_{18})^{(2)}(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}G_{16}^* \right) \end{aligned}$$

$$\begin{aligned}
& \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(18)}T_{17}^* + (b_{17})^{(2)}s_{(16),(18)}T_{16}^* \right) \} = 0 \\
& + \\
& \left( (\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \\
& \left[ ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)})(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(q_{20})^{(3)}G_{20}^* \right] \\
& \left( ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(21)}T_{21}^* + (b_{21})^{(3)}s_{(20),(21)}T_{21}^* \right) \\
& + \left( ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)})(q_{20})^{(3)}G_{20}^* + (a_{20})^{(3)}(q_{21})^{(1)}G_{21}^* \right) \\
& \left( ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(20)}T_{21}^* + (b_{21})^{(3)}s_{(20),(20)}T_{20}^* \right) \\
& \left( ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) \\
& \left( ((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \right) \\
& + \left( ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \right) (q_{22})^{(3)}G_{22} \\
& + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)}(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G_{20}^*) \\
& \left( ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(22)}T_{21}^* + (b_{21})^{(3)}s_{(20),(22)}T_{20}^* \right) \} = 0 \\
& + \\
& \left( (\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \right) \{ (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \\
& \left[ ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(q_{24})^{(4)}G_{24}^* \right] \\
& \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(25)}T_{25}^* + (b_{25})^{(4)}s_{(24),(25)}T_{25}^* \right) \\
& + \left( ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)})(q_{24})^{(4)}G_{24}^* + (a_{24})^{(4)}(q_{25})^{(4)}G_{25}^* \right) \\
& \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(24)}T_{25}^* + (b_{25})^{(4)}s_{(24),(24)}T_{24}^* \right) \\
& \left( ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
& \left( ((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
& + \left( ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) (q_{26})^{(4)}G_{26} \\
& + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)}(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G_{24}^*) \\
& \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(26)}T_{25}^* + (b_{25})^{(4)}s_{(24),(26)}T_{24}^* \right) \} = 0 \\
& + \\
& \left( (\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \\
& \left[ ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)}G_{28}^* \right] \\
& \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(29)}T_{29}^* + (b_{29})^{(5)}s_{(28),(29)}T_{29}^* \right) \\
& + \left( ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)}G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)}G_{29}^* \right) \\
& \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(28)}T_{29}^* + (b_{29})^{(5)}s_{(28),(28)}T_{28}^* \right) \\
& \left( ((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
& \left( ((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right) \\
& + \left( ((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)}G_{30} \\
& + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)}(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G_{28}^*) \\
& \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(30)}T_{29}^* + (b_{29})^{(5)}s_{(28),(30)}T_{28}^* \right) \} = 0 \\
& + \\
& \left( (\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \{ (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \\
& \left[ ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)}G_{32}^* \right] \\
& \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T_{33}^* + (b_{33})^{(6)}s_{(32),(33)}T_{33}^* \right) \\
& + \left( ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G_{32}^* + (a_{32})^{(6)}(q_{33})^{(6)}G_{33}^* \right) \\
& \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T_{33}^* + (b_{33})^{(6)}s_{(32),(32)}T_{32}^* \right) \\
& \left( ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
& \left( ((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
& + \left( ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) (q_{34})^{(6)}G_{34} \\
& + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)}(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G_{32}^*)
\end{aligned}$$

$$\left\{ \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right\} = 0$$

+

$$\begin{aligned} & \left( (\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \left\{ \left( (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \right) \right. \\ & \left. \left[ \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \right\} \\ & \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \\ & + \left( (\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)} \right) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \\ & \left( (\lambda)^{(7)} \right)^2 + \left( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} \\ & \left( (\lambda)^{(7)} \right)^2 + \left( (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} \right) (\lambda)^{(7)} \\ & + \left( (\lambda)^{(7)} \right)^2 + \left( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} \right) (\lambda)^{(7)} (q_{38})^{(7)} G_{38} \\ & + \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left( (a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left. \left( (\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)} \right) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right\} = 0 \end{aligned}$$

## REFERENCES

- =====
- =====
- (1) A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80<sup>th</sup> birthday
  - (2) FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188
  - (3) HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [<http://www.eolss.net>]
  - (4) MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097
  - (5) STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-1006
  - (6) FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010
  - (7) HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [<http://www.eolss.net>]
  - (8) MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097
  - (8A) STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-1006
  - (8B) FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010

(9)<sup>a b c</sup> Einstein, A. (1905), "Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?", *Annalen der Physik* **18**: 639 Bibcode 1905AnP...323..639E, DOI:10.1002/andp.19053231314. See also the English translation.

(10)<sup>a b</sup> Paul Allen Tipler, Ralph A. Llewellyn (2003-01), *Modern Physics*, W. H. Freeman and Company, pp. 87–88, ISBN 0-7167-4345-0

(11)<sup>a b</sup> Rainville, S. et al. World Year of Physics: A direct test of  $E=mc^2$ . *Nature* 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.

(12)<sup>^</sup> In F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy

(13)<sup>^</sup> Note that the relativistic mass, in contrast to the rest mass  $m_0$ , is not a relativistic invariant, and that the velocity is not a Minkowski four-vector, in contrast to the quantity  $dx^\mu/d\tau$ , where  $\tau$  is the differential of the proper time. However, the energy-momentum four-vector is a genuine Minkowski four-vector, and the intrinsic origin of the square-root in the definition of the relativistic mass is the distinction between  $d\mathbf{x}$  and  $d\mathbf{t}$ .

(14)<sup>^</sup> Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0

(15)<sup>^</sup> Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8

(16)<sup>^</sup> Hans, H. S.; Puri, S. P. (2003). *Mechanics* (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433

(17)<sup>^</sup> E. F. Taylor and J. A. Wheeler, **Spacetime Physics**, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.

(18)<sup>^</sup> Mould, Richard A. (2002). *Basic relativity* (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126

(19)<sup>^</sup> Chow, Tail L. (2006). *Introduction to electromagnetic theory: a modern perspective*. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392

(20)<sup>^</sup> [2] Cockcroft-Walton experiment

(21)<sup>a b c</sup> Conversions used: 1956 International (Steam) Table (IT) values where one calorie  $\equiv$  4.1868 J and one BTU  $\equiv$  1055.05585262 J. Weapons designers' conversion value of one gram TNT  $\equiv$  1000 calories used.

(22)<sup>^</sup> Assuming the dam is generating at its peak capacity of 6,809 MW.

(23)<sup>^</sup> Assuming a 90/10 alloy of Pt/Ir by weight, a  $C_p$  of 25.9 for Pt and 25.1 for Ir, a Pt-dominated average  $C_p$  of 25.8, 5.134 moles of metal, and 132 J.K<sup>-1</sup> for the prototype. A variation of  $\pm 1.5$  picograms is of course, much smaller than the actual uncertainty in the mass of the international prototype, which are  $\pm 2$  micrograms.

(24)<sup>^</sup> [3] Article on Earth rotation energy. Divided by  $c^2$ .

(25)<sup>a b</sup> Earth's gravitational self-energy is  $4.6 \times 10^{10}$  that of Earth's total mass, or 2.7 trillion metric tons. Citation: *The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO)*, T. W. Murphy, Jr. et al. University of Washington, Dept. of Physics (132 kB PDF, here.).

(26)<sup>^</sup> There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be *minimal coupling*, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.

- (27)<sup>△</sup> G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Physical Review D* 14:3432–3450 (1976).
- (28)<sup>△</sup> A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", *Physics Letters* 59B:85 (1975).
- (29)<sup>△</sup> F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", *Physical Review D* 30:2212.
- (30)<sup>△</sup> Rubakov V. A. "Monopole Catalysis of Proton Decay", *Reports on Progress in Physics* 51:189–241 (1988).
- (31)<sup>△</sup> S.W. Hawking "Black Holes Explosions?" *Nature* 248:30 (1974).
- (32)<sup>△</sup> Einstein, A. (1905), "Zur Elektrodynamik bewegter Körper." (PDF), *Annalen der Physik* 17: 891–921, Bibcode 1905AnP...322...891E, DOI:10.1002/andp.19053221004. English translation.
- (33)<sup>△</sup> See e.g. Lev B.Okun, *The concept of Mass*, *Physics Today* 42 (6), June 1969, p. 31–36, [http://www.physicstoday.org/vol-42/iss-6/vol42no6p31\\_36.pdf](http://www.physicstoday.org/vol-42/iss-6/vol42no6p31_36.pdf)
- (34)<sup>△</sup> Max Jammer (1999), *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, p. 51, ISBN 0-691-01017-X
- (35)<sup>△</sup> Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", *Foundations of Physics* (Springer) 6: 115–124, Bibcode 1976FoPh....6..115E, DOI:10.1007/BF00708670
- (36)<sup>△ a b</sup> Janssen, M., Mecklenburg, M. (2007), *From classical to relativistic mechanics: Electromagnetic models of the electron.*, in V. F. Hendricks, et al., *Interactions: Mathematics, Physics and Philosophy* (Dordrecht: Springer): 65–134
- (37)<sup>△ a b</sup> Whittaker, E.T. (1951–1953), 2. Edition: *A History of the theories of aether and electricity, vol. 1: The classical theories / vol. 2: The modern theories 1900–1926*, London: Nelson
- (38)<sup>△</sup> Miller, Arthur I. (1981), *Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911)*, Reading: Addison–Wesley, ISBN 0-201-04679-2
- (39)<sup>△ a b</sup> Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), *Séminaire Poincaré* 1: 1–22
- (40)<sup>△</sup> Philip Ball (Aug 23, 2011). "Did Einstein discover  $E = mc^2$ ?" *Physics World*.
- (41)<sup>△</sup> Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* 42 (8): 540–543, DOI:10.1364/JOSA.42.000540
- (42)<sup>△</sup> Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.
- (43)<sup>△</sup> Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* 50 (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764
- (44)<sup>△</sup> Ohanian, Hans (2008), "Did Einstein prove  $E=mc^2$ ?", *Studies In History and Philosophy of Science Part B* 40 (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002
- (45)<sup>△</sup> Hecht, Eugene (2011), "How Einstein confirmed  $E_0=mc^2$ ", *American Journal of Physics* 79 (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223
- (46)<sup>△</sup> Rohrlich, Fritz (1990), "An elementary derivation of  $E=mc^2$ ", *American Journal of Physics* 58 (4): 348–

(47) (1996). *Lise Meitner: A Life in Physics*. California Studies in the History of Science. **13**. Berkeley: University of California Press. pp. 236–237. ISBN 0-520-20860-

(48)^ UIBK.ac.at

(49)^ J. J. L. Morton; *et al.* (2008). "Solid-state quantum memory using the  $^{31}\text{P}$  nuclear spin". *Nature* **455** (7216): 1085–1088. Bibcode 2008Natur.455.1085M.DOI:10.1038/nature07295.

(50)^ S. Weisner (1983). "Conjugate coding". *Association of Computing Machinery, Special Interest Group in Algorithms and Computation Theory* **15**: 78–88.

(51)^ A. Zeilinger, *Dance of the Photons: From Einstein to Quantum Teleportation*, Farrar, Straus & Giroux, New York, 2010, pp. 189, 192, ISBN 0374239665

(52)^ B. Schumacher (1995). "Quantum coding". *Physical Review A* **51** (4): 2738–2747. Bibcode 1995PhRvA..51.2738S. DOI:10.1103/PhysRevA.51.2738.

(53)Delamotte, Bertrand; *A hint of renormalization*, American Journal of Physics 72 (2004) pp. 170–184. Beautiful elementary introduction to the ideas, no prior knowledge of field theory being necessary. Full text available at: [hep-th/0212049](http://hep-th/0212049)

(54)Baez, John; *Renormalization Made Easy*, (2005). A qualitative introduction to the subject.

(55)Blechman, Andrew E. ; *Renormalization: Our Greatly Misunderstood Friend*, (2002). Summary of a lecture; has more information about specific regularization and divergence-subtraction schemes.

(56)Cao, Tian Yu & Schweber, Silvian S. ; *The Conceptual Foundations and the Philosophical Aspects of Renormalization Theory*, Synthese, 97(1) (1993), 33–108.

(57)Shirkov, Dmitry; *Fifty Years of the Renormalization Group*, C.E.R.N. Courier 41(7) (2001). Full text available at: *I.O.P Magazines*.

(58)E. Elizalde; *Zeta regularization techniques with Applications*.

(59)N. N. Bogoliubov, D. V. Shirkov (1959): *The Theory of Quantized Fields*. New York, Interscience. The first text-book on the renormalization group theory.

(60)Ryder, Lewis H. ; *Quantum Field Theory* (Cambridge University Press, 1985), ISBN 0-521-33859-X Highly readable textbook, certainly the best introduction to relativistic Q.F.T. for particle physics.

(61)Zee, Anthony; *Quantum Field Theory in a Nutshell*, Princeton University Press (2003) ISBN 0-691-01019-6. Another excellent textbook on Q.F.T.

(62)Weinberg, Steven; *The Quantum Theory of Fields* (3 volumes) Cambridge University Press (1995). A monumental treatise on Q.F.T. written by a leading expert, *Nobel laureate 1979*.

(63)Pokorski, Stefan; *Gauge Field Theories*, Cambridge University Press (1987) ISBN 0-521-47816-2.

(64)'t Hooft, Gerard; *The Glorious Days of Physics – Renormalization of Gauge theories*, lecture given at Erice (August/September 1998) by the *Nobel laureate 1999* . Full text available at: [hep-th/9812203](http://hep-th/9812203).

(65)Rivasseau, Vincent; *An introduction to renormalization*, Poincaré Seminar (Paris, Oct. 12, 2002), published in: Duplantier, Bertrand; Rivasseau, Vincent (Eds.) ; *Poincaré Seminar 2002*, Progress in Mathematical Physics



- (66) Rivasseau, Vincent; *From perturbative to constructive renormalization*, Princeton University Press (1991) ISBN 0-691-08530-7. Full text available in *PostScript*.
- (67) H.P. Nilles, Phys. Rep. 110, 1 (1984);
- (68) H.E. Haber and G.L. Kane, Phys. Rep. 117, 75 (1985);
- (69) R. Barbieri, Riv. Nuov. Cim. 11, 1 (1988).
- (70) J.F. Gunion and H.E. Haber, Nucl. Phys. B272, 1 (1986); B402, 567(E) (1993).
- (71) A. Sirlin, Nucl. Phys. B71, 29 (1974); Rev. Mod. Phys. 50, 573 (1978);
- (72) W.J. Marciano and A. Sirlin, Nucl. Phys. B93, 303 (1975).
- (73) A. Denner and T. Sack, Nucl. Phys. B347, 203 (1990).
- (74) B.A. Kniehl and A. Pilaftsis, Nucl. Phys. B474, 286 (1996).
- (75) J. Guasch, J. Sol`a, and W. Hollik, Phys. Lett. B 437, 88 (1998);
- (76) H. Eberl, S. Kraml, and W. Majerotto, JHEP 9905, 016 (1999);
- (77) J. Guasch, W. Hollik, and J. Sol`a, Phys. Lett. B 510, 211 (2001).
- [78] P. Gambino, P.A. Grassi, and F. Madricardo, Phys. Lett. B 454, 98 (1999).
- (79) B.A. Kniehl, F. Madricardo, and M. Steinhauser, Phys. Rev. D 62, 073010 (2000).
- (80) A. Barroso, L. Br`ucher, and R. Santos, Phys. Rev. D 62, 096003 (2000).
- (81) N. Cabibbo, Phys. Rev. Lett. 19, 531 (1963);
- (82) M. Kobayashi and M. Maskawa, Prog. Theor. Phys. 49, 652 (1973).
- (83) J.M. Cornwall, in Proceedings of the French-American Seminar on Theoretical Aspects of Quantum Chromodynamics, Marseille, France, 1981, edited by J.W. Dash (Centre de Physique Th´eorique, Marseille, 1982);
- (84) J.M. Cornwall, Phys. Rev. D 26, 1453 (1982);
- (85) J.M. Cornwall and J. Papavassiliou, Phys. Rev. D 40, 3474 (1989).

---

#### Acknowledgments:

The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's Letters, Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive

**First Author:** <sup>1</sup>Mr. K. N.Prasanna Kumar has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding **Author:drknpkumar@gmail.com**

**Second Author:** <sup>2</sup>Prof. B.S Kiranagi is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

**Third Author:** <sup>3</sup>Prof. C.S. Bagewadi is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India

---

