

## Theory of Alfa Ray Production, Quantum Tunneling, Redundancy, Entropy, Event, Cause, Space, Time, Storage Ability and Entanglement-A Tarantula-Guillemot Model

<sup>1</sup>Dr. K. N. Prasanna Kumar, <sup>2</sup>Prof. B. S. Kiranagi, <sup>3</sup>Prof. C. S. Bagewadi

**ABSTRACT:** We propose a Theory of Alpha ray production(Radioactive decay), Quantum Tunneling, Redundancy, Entropy, Event, Cause, Space, Time, Storage ability, Quantum entanglement, Noise, Errors, by a Model that concatenates and consummates these important variables and provide predictive capacity, Stability analysis, and Solutional behaviour of the system,

### I. INTRODUCTION

What is an event? Or for that matter an ideal event? An event is a singularity or rather a set of singularities or set of singular points characterizing a mathematical curve, a physical state of affairs, a psychological person or a moral person. Singularities are turning points and points of inflection: they are bottle necks, foyers and centers; they are points of fusion; condensation and boiling; points of tears and joy; sickness and health; hope and anxiety; they are so to say "sensitive" points; such singularities should not be confused or confounded, aggravated or exacerbated with personality of a system expressing itself; or the individuality and idiosyncrasies of a system which is designated with a proposition. They should also not be fused with the generalizational concept or universalistic axiomatic predications and postulation alcovishness, or the dipsomaniac flageolet dirge of a concept. Possible a concept could be signified by a figurative representation or a schematic configuration. "Singularity is essentially, pre individual, and has no personalized bias in it, or for that matter a prejudice or pre circumspection of a conceptual scheme. It is in this sense we can define a "singularity" as being neither affirmative nor non affirmative. it can be positive or negative; it can create or destroy. On the other hand it must be noted that singularity is different both in its thematic discursive from the run of the mill day to day musings and mundane drooling. There are in that sense "extra-ordinary".

Each singularity is a source and resource, the origin, reason and raison d'être of a mathematical series, it could be any series any type, and that is interpolated or extrapolated to the structural location of the destination of another singularity. This according to this standpoint, there are different, multifarious, myriad, series in a structure. In the eventuality of the fact that we conduct an unbiased and prudent examination of the series belonging to different "singularities" we can come to indubitable conclusion that the "singularity" of one system is different from the "other system" in the subterranean realm and ceratoid dualism of comparison and contrast

EPR experiment derived that there exists a communications between two particles. We go a further step to say that there exists a channel of communication however slovenly, inept, clumpy, between the two singularities. It is also possible the communication exchange could be one of belligerence, cantankerousness, tempestuousness, astutely truculent, with ensorcelled frenzy. That does not matter. All we are telling is that singularities communicate with each other.

Now, how do find the reaction of systems to these singularities. You do the same thing a boss does for you. "Problematize" the events and see how you behave. I will resort to "pressure tactics". "intimidation of deriding report", or "cut in the increment" to make you undergo trials, travails and tribulations. i am happy to see if you improve your work; but may or may not be sad if you succumb to it and hang yourself! We do the same thing with systems. systems show conducive response, felicitous reciprocation or behave erratically with inner roil, eponymous radicalism without and with blitz conviction say like a solipsist nature of bellicose and blustering particles, or for that matter coruscation, trepidational motion in fluid flows, or seemingly perfidious incendiaries in gormandizing fellow elementary particles, abnormal ebullitions, surcharges calumniation and unwarranted (you think so but the system does not!) unrighteous fulminations.

So the point that is made here is " like we problematize the "events" to understand the human behaviour we have to "problematize" the events of systems to understand their behaviour.

This statement is made in connection to the fact that there shall be creation or destruction of particles or complete obliteration of the systems (blackhole evaporation) or obfuscation of results. Some systems are like "inside traders" they will not put signature at all! How do you find they did it! Anyway, there are possibilities of a CIA/CBI finding out as they recently did! So we can do the same thing with systems to. This is accentuation, corroboration, fortification, .commendatory note to explain the various coefficients we have used in the model as also the dissipations called for In the bank example we have clarified that various systems are individually conservative, and their conservativeness extends holistically too.that one law is universal does not mean there is complete adjudication of nonexistence of totality or global or holistic figure. Total always exists and "individual" systems always exist, if we do not bring Kant in to picture! For the time being let us not! Equations would become more eneuretic and frenzied..

We TAKE In to consideration the following parameter:

- (1) Alpha ray production(Radioactive decay)
- (2) Quantum Tunneling
- (3) Redundancy
- (4) Entropy
- (5) Event
- (6) Cause
- (7) Space
- (8) Time
- (9) Storage ability
- (10)Quantum entanglement
- (11)Noise Errors.

**QUANTUM TUNNELING AND ALPHA RAY PRODUCTION(RADIOACTIVE DECAY)**

**MODULE NUMBERED ONE.**

**NOTATION:**

- $G_{13}$  : CATEGORY ONE OF QUANTUM TUNNELING  
 $G_{14}$  : CATEGORY TWO OF QUANTUM TUNNELING  
 $G_{15}$  : CATEGORY THREE OF QUANTUM TUNNELING  
 $T_{13}$  : CATEGORY ONE OF ALPHA RAY PRODUCTION(RADIOACTIVE DECAY)  
 $T_{14}$  : CATEGORY TWO OF ALPHA RAY PRODUCTION (RADIOACTIVE DECAY)  
 $T_{15}$  :CATEGORY THREE OF ALPHA RAY PRODUCTION(RADIOACTIVE DECAY)
- 

**ENTROPY AND REDUNDANCY**  
**MODULE NUMBERED TWO:**

- $G_{16}$  : CATEGORY ONE OF ENTROPY  
 $G_{17}$  : CATEGORY TWO OF ENTROPY  
 $G_{18}$  : CATEGORY THREE OF ENTROPY  
 $T_{16}$  :CATEGORY ONE OF REDUNDANCY  
 $T_{17}$  : CATEGORY TWO OF REDUNDANCY  
 $T_{18}$  : CATEGORY THREE OF REDUNDANCY
- 

**NOISE AND ERROR IN QUANTUM COMPUTATION:**  
**MODULE NUMBERED THREE:**

- $G_{20}$  : CATEGORY ONE OF ERRORS  
 $G_{21}$  :CATEGORY TWO OF ERRORS  
 $G_{22}$  : CATEGORY THREE OF ERRORS  
 $T_{20}$  : CATEGORY ONE OF NOISE  
 $T_{21}$  :CATEGORY TWO OF NOISE  
 $T_{22}$  : CATEGORY THREE OF NOISE
- 

**CAUSE AND EVENT:**  
**MODULE NUMBERED FOUR:**

- $G_{24}$  : CATEGORY ONE OF CAUSE  
 $G_{25}$  : CATEGORY TWO OF CAUSE  
 $G_{26}$  : CATEGORY THREE OF CAUSE  
 $T_{24}$  :CATEGORY ONE OF EVENT  
 $T_{25}$  :CATEGORY TWO OF EVENT  
 $T_{26}$  : CATEGORY THREE OF EVENT
- 

**SPACE AND TIME:**  
**MODULE NUMBERED FIVE:**

- $G_{28}$  : CATEGORY ONE OF TIME  
 $G_{29}$  : CATEGORY TWO OF TIME  
 $G_{30}$  :CATEGORY THREE OF TIME  
 $T_{28}$  :CATEGORY ONE OF SPACE  
 $T_{29}$  :CATEGORY TWO OF SPACE  
 $T_{30}$  :CATEGORY THREE OF SPACE
-

**STORAGE ABILITY AND ENTANGLEMENT :**  
**MODULE NUMBERED SIX:**

$G_{32}$  : CATEGORY ONE OF ENTANGLEMENT  
 $G_{33}$  : CATEGORY TWO OF ENTANGLEMENT  
 $G_{34}$  : CATEGORY THREE OF ENTANGLEMENT  
 $T_{32}$  : CATEGORY ONE OF STORAGE ABILITY  
 $T_{33}$  : CATEGORY TWO OF STORAGE ABILITY  
 $T_{34}$  : CATEGORY THREE OF STORAGE ABILITY

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$ :  
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$   
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}$ ,  
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$   
 $, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$   
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}$ ,  
 $(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

are Dissipation coefficients.

**QUANTUM TUNNELING AND ALPHA RAY PRODUCTION(RADIOACTIVE DECAY)**  
**MODULE NUMBERED ONE**

The differential system of this model is now (Module Numbered one).

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} . \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} . \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} . \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} . \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} . \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} . \\ + (a''_{13})^{(1)}(T_{14}, t) &= \text{First augmentation factor} . \\ - (b''_{13})^{(1)}(G, t) &= \text{First detritions factor} . \end{aligned}$$

**ENTROPY AND REDUNDANCY**  
**MODULE NUMBERED TWO**

The differential system of this model is now ( Module numbered two).

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} . \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} . \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} . \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} . \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} . \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} . \\ + (a''_{16})^{(2)}(T_{17}, t) &= \text{First augmentation factor} . \\ - (b''_{16})^{(2)}((G_{19}), t) &= \text{First detritions factor} . \end{aligned}$$

**NOISE AND ERROR IN QUANTUM COMPUTATION:**  
**MODULE NUMBERED THREE**

The differential system of this model is now (Module numbered three).

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} . \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} . \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} . \end{aligned}$$

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \cdot \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \cdot \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \cdot \\ + (a''_{20})^{(3)}(T_{21}, t) &= \text{First augmentation factor.} \\ - (b_{20})^{(3)}(G_{23}, t) &= \text{First detritions factor.} \end{aligned}$$

**CAUSE AND EVENT:**

**MODULE NUMBERED FOUR:**

The differential system of this model is now (Module numbered Four).

$$\begin{aligned} \frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \cdot \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \cdot \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \cdot \\ \frac{dT_{24}}{dt} &= (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \cdot \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \cdot \\ \frac{dT_{26}}{dt} &= (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \cdot \\ + (a''_{24})^{(4)}(T_{25}, t) &= \text{First augmentation factor.} \\ - (b''_{24})^{(4)}((G_{27}), t) &= \text{First detritions factor.} \end{aligned}$$

**SPACE AND TIME:**

**MODULE NUMBERED FIVE**

The differential system of this model is now (Module number five).

$$\begin{aligned} \frac{dG_{28}}{dt} &= (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \cdot \\ \frac{dG_{29}}{dt} &= (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \cdot \\ \frac{dG_{30}}{dt} &= (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \cdot \\ \frac{dT_{28}}{dt} &= (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \cdot \\ \frac{dT_{29}}{dt} &= (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \cdot \\ \frac{dT_{30}}{dt} &= (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \cdot \\ + (a''_{28})^{(5)}(T_{29}, t) &= \text{First augmentation factor.} \\ - (b''_{28})^{(5)}((G_{31}), t) &= \text{First detritions factor.} \end{aligned}$$

**STORAGE ABILITY AND ENTANGLEMENT :**  
**MODULE NUMBERED SIX**

The differential system of this model is now (Module numbered Six).

$$\begin{aligned} \frac{dG_{32}}{dt} &= (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \cdot \\ \frac{dG_{33}}{dt} &= (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \cdot \\ \frac{dG_{34}}{dt} &= (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \cdot \\ \frac{dT_{32}}{dt} &= (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \cdot \\ \frac{dT_{33}}{dt} &= (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \cdot \\ \frac{dT_{34}}{dt} &= (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \cdot \\ + (a''_{32})^{(6)}(T_{33}, t) &= \text{First augmentation factor.} \\ - (b''_{32})^{(6)}((G_{35}), t) &= \text{First detritions factor.} \end{aligned}$$

**HOLISTIC CONCATENATE SYTEMAL EQUATIONS HENCEFORTH REFERRED TO AS “GLOBAL EQUATIONS”**

- (1) Alpha ray production(Radioactive decay)
- (2) Quantum Tunneling
- (3) Redundancy
- (4) Entropy
- (5) Event
- (6) Cause
- (7) Space
- (8) Time
- (9) Storage ability
- (10) Quantum entanglement
- (11) Noise
- (12) Errors

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - \left[ \begin{array}{|c|c|c|} \hline (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14}, t) & +(a''_{16})^{(2,2)}(T_{17}, t) & +(a''_{20})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{13} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline +(a''_{24})^{(4,4,4,4)}(T_{25}, t) & +(a''_{28})^{(5,5,5,5)}(T_{29}, t) & +(a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - \left[ \begin{array}{|c|c|c|} \hline (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14}, t) & +(a''_{17})^{(2,2)}(T_{17}, t) & +(a''_{21})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{14} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline +(a''_{25})^{(4,4,4,4)}(T_{25}, t) & +(a''_{29})^{(5,5,5,5)}(T_{29}, t) & +(a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - \left[ \begin{array}{|c|c|c|} \hline (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14}, t) & +(a''_{18})^{(2,2)}(T_{17}, t) & +(a''_{22})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{15} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline +(a''_{26})^{(4,4,4,4)}(T_{25}, t) & +(a''_{30})^{(5,5,5,5)}(T_{29}, t) & +(a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] \end{aligned}$$

Where  $(a'_{13})^{(1)}(T_{14}, t)$ ,  $(a'_{14})^{(1)}(T_{14}, t)$ ,  $(a'_{15})^{(1)}(T_{14}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 $(a''_{16})^{(2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3  
 $(a''_{20})^{(3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3  
 $(a''_{24})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3  
 $(a''_{28})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3  
 $(a''_{32})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3.

$$\begin{aligned} \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - \left[ \begin{array}{|c|c|c|} \hline (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G, t) & -(b''_{16})^{(2,2)}(G_{19}, t) & -(b''_{20})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{13} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline -(b''_{24})^{(4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ \hline \end{array} \right] \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - \left[ \begin{array}{|c|c|c|} \hline (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G, t) & -(b''_{17})^{(2,2)}(G_{19}, t) & -(b''_{21})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{14} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline -(b''_{25})^{(4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ \hline \end{array} \right] \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - \left[ \begin{array}{|c|c|c|} \hline (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G, t) & -(b''_{18})^{(2,2)}(G_{19}, t) & -(b''_{22})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{15} \cdot \\ & \left[ \begin{array}{|c|c|c|} \hline -(b''_{26})^{(4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6)}(G_{35}, t) \\ \hline \end{array} \right] \end{aligned}$$

Where  $-(b''_{13})^{(1)}(G, t)$ ,  $-(b''_{14})^{(1)}(G, t)$ ,  $-(b''_{15})^{(1)}(G, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{16})^{(2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{20})^{(3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a'_{13})^{(1,1)}(T_{14}, t) & + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{16} \cdot \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - \left[ \begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a'_{14})^{(1,1)}(T_{14}, t) & + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{17} \cdot \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - \left[ \begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a'_{15})^{(1,1)}(T_{14}, t) & + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{18} \cdot \end{aligned}$$

Where  $+(a'_{16})^{(2)}(T_{17}, t)$ ,  $+(a'_{17})^{(2)}(T_{17}, t)$ ,  $+(a'_{18})^{(2)}(T_{17}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 $+(a'_{13})^{(1,1)}(T_{14}, t)$ ,  $+(a'_{14})^{(1,1)}(T_{14}, t)$ ,  $+(a'_{15})^{(1,1)}(T_{14}, t)$  are second augmentation coefficient for category 1, 2 and 3  
 $+(a'_{20})^{(3,3,3)}(T_{21}, t)$ ,  $+(a'_{21})^{(3,3,3)}(T_{21}, t)$ ,  $+(a'_{22})^{(3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3 .

$$\begin{aligned} \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - \left[ \begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b'_{13})^{(1,1)}(G, t) & - (b'_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{16} \cdot \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - \left[ \begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b'_{14})^{(1,1)}(G, t) & - (b'_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{17} \cdot \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - \left[ \begin{array}{ccc} (b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}, t) & - (b'_{15})^{(1,1)}(G, t) & - (b'_{22})^{(3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \end{array} \right] T_{18} \cdot \end{aligned}$$

where  $-(b''_{16})^{(2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2)}(G_{19}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1)}(G, t)$  are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1,2 and 3

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) & + (a'_{16})^{(2,2,2)}(T_{17}, t) & + (a'_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{20} \cdot \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - \left[ \begin{array}{ccc} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) & + (a'_{17})^{(2,2,2)}(T_{17}, t) & + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{21} \cdot \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \end{array} \right] G_{22} \cdot \end{aligned}$$

$\boxed{+(a''_{20})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3)}(T_{21}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)}$  are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - \left[ \begin{array}{ccc} \boxed{(b'_{20})^{(3)}} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{20} \cdot \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - \left[ \begin{array}{ccc} \boxed{(b'_{21})^{(3)}} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{21} \cdot \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - \left[ \begin{array}{ccc} \boxed{(b'_{22})^{(3)}} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{22} \cdot \end{aligned}$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{24}}{dt} &= (a_{24})^{(4)}G_{25} - \left[ \begin{array}{ccc} \boxed{(a'_{24})^{(4)}} \boxed{+(a''_{24})^{(4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{24} \cdot \\ \frac{dG_{25}}{dt} &= (a_{25})^{(4)}G_{24} - \left[ \begin{array}{ccc} \boxed{(a'_{25})^{(4)}} \boxed{+(a''_{25})^{(4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{25} \cdot \\ \frac{dG_{26}}{dt} &= (a_{26})^{(4)}G_{25} - \left[ \begin{array}{ccc} \boxed{(a'_{26})^{(4)}} \boxed{+(a''_{26})^{(4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{26} \cdot \end{aligned}$$

Where  $\boxed{(a''_{24})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{25})^{(4)}(T_{25}, t)}$ ,  $\boxed{(a''_{26})^{(4)}(T_{25}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5)}(T_{29}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, and 3.

$$\begin{aligned} \frac{dT_{24}}{dt} &= (b_{24})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{24})^{(4)} \boxed{-(b''_{24})^{(4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{24} \cdot \\ \frac{dT_{25}}{dt} &= (b_{25})^{(4)}T_{24} - \left[ \begin{array}{ccc} (b'_{25})^{(4)} \boxed{-(b''_{25})^{(4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{25} \cdot \\ \frac{dT_{26}}{dt} &= (b_{26})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{26})^{(4)} \boxed{-(b''_{26})^{(4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{26} \cdot \end{aligned}$$

Where  $\boxed{-(b''_{24})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4)}(G_{27}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{13})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1)}(G, t)}$   
 are fourth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)}$   
 are fifth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$   
 are sixth detrition coefficients for category 1, 2 and 3 .

$$\begin{aligned} \frac{dG_{28}}{dt} &= (a_{28})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{28})^{(5)} \boxed{+(a''_{28})^{(5)}(T_{29}, t)} & \boxed{+(a''_{24})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{28} \cdot \\ \frac{dG_{29}}{dt} &= (a_{29})^{(5)}G_{28} - \left[ \begin{array}{ccc} (a'_{29})^{(5)} \boxed{+(a''_{29})^{(5)}(T_{29}, t)} & \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{29} \cdot \\ \frac{dG_{30}}{dt} &= (a_{30})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{30})^{(5)} \boxed{+(a''_{30})^{(5)}(T_{29}, t)} & \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \end{array} \right] G_{30} \cdot \end{aligned}$$

Where  $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 And  $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$  are second augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3  
 $\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3  
 $\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, 3 .

$$\begin{aligned} \frac{dT_{28}}{dt} &= (b_{28})^{(5)}T_{29} - \left[ \begin{array}{ccc} (b'_{28})^{(5)} \boxed{-(b''_{28})^{(5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{28} \cdot \\ \frac{dT_{29}}{dt} &= (b_{29})^{(5)}T_{28} - \left[ \begin{array}{ccc} (b'_{29})^{(5)} \boxed{-(b''_{29})^{(5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{29} \cdot \\ \frac{dT_{30}}{dt} &= (b_{30})^{(5)}T_{29} - \left[ \begin{array}{ccc} (b'_{30})^{(5)} \boxed{-(b''_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \end{array} \right] T_{30} \cdot \end{aligned}$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1, 2 and 3



$-(b''_{32})^{(6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6)}(G_{35}, t)$  are third detrition coefficients for category 1,2 and 3  
 $-(b''_{13})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1,2, and 3  
 $-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1,2, and 3  
 $-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1,2, and 3.

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a''_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{32} .$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{ccc} (a''_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{33} .$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a''_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{34} .$$

$+(a''_{32})^{(6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6)}(T_{33}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 $+(a''_{28})^{(5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5)}(T_{29}, t)$  are second augmentation coefficients for category 1, 2 and 3  
 $+(a''_{24})^{(4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4)}(T_{25}, t)$  are third augmentation coefficients for category 1, 2 and 3  
 $+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$  - are fourth augmentation coefficients  
 $+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$  - fifth augmentation coefficients  
 $+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$  sixth augmentation coefficients .

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{32} .$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{33} .$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{34} .$$

$-(b''_{32})^{(6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6)}(G_{35}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4)}(G_{27}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2, and 3  
 $-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2, and 3  
 $-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2, and 3

Where we suppose.

- (A)  $(a_i)^{(1)}, (a_i'')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0$ ,  
 $i, j = 13, 14, 15$   
 (B) The functions  $(a_i'')^{(1)}, (b_i'')^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (\tau_i)^{(1)}$ :

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}.$$

$$(C) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13, 14, 15$ .

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(M_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(M_{13})^{(1)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions

$(a_i'')^{(1)}(T'_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the first augmentation coefficient WOULD be absolutely continuous. .

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

(D)  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1.$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

(E) There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together with  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1.$$

Where we suppose.

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18.$$

The functions  $(a_i'')^{(2)}, (b_i'')^{(2)}$  are positive continuous increasing and bounded..

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$  :

$$(a_i'')^{(2)}(T'_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)}.$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)}.$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)}.$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)}.$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  :

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$ .

They satisfy Lipschitz condition:.

$$|(a_i'')^{(2)}(T'_{17}, t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T'_{17} - T_{17}| e^{-(M_{16})^{(2)}t}$$

$$|(b_i'')^{(2)}(G'_{19}, t) - (b_i'')^{(2)}(G_{19}, t)| < (\hat{k}_{16})^{(2)} \|G_{19} - G'_{19}\| e^{-(M_{16})^{(2)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(2)}(T'_{17}, t)$  and  $(a_i'')^{(2)}(T_{17}, t)$ .  $(T'_{17}, t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a_i'')^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a_i'')^{(2)}(T_{17}, t)$ , the SECOND augmentation coefficient would be absolutely continuous. .

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  :

(F)  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1.$$

**Definition of**  $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$ , satisfy the inequalities .

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1.$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1.$$

Where we suppose.

(G)  $(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22$

The functions  $(a''_i)^{(3)}, (b''_i)^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}.$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$ :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$ .

They satisfy Lipschitz condition:

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T'_{21} - T_{21}| e^{-(M_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G'_{23} - G_{23}'|| e^{-(M_{20})^{(3)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(3)}(T'_{21}, t)$  and  $(a''_i)^{(3)}(T_{21}, t)$ .  $(T'_{21}, t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a''_i)^{(3)}(T'_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a''_i)^{(3)}(T'_{21}, t)$ , the THIRD augmentation coefficient, would be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ :

(H)  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1.$$

There exists two constants There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1.$$

Where we suppose.

(I)  $(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26$

(J) The functions  $(a''_i)^{(4)}, (b''_i)^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}.$$

$$(K) \quad \lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$ :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$ .

They satisfy Lipschitz condition:

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(M_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G'_{27}), t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G'_{27}) - (G_{27})'|| e^{-(M_{24})^{(4)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(4)}(T'_{25}, t)$  and  $(a''_i)^{(4)}(T_{25}, t)$ .  $(T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a''_i)^{(4)}(T'_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 4$  then the function  $(a''_i)^{(4)}(T'_{25}, t)$ , the FOURTH augmentation coefficient WOULD be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ :

(L)  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

(M) 
$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1.$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$ :

(N) There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}$ ,  $(\hat{k}_{24})^{(4)}$ ,  $(\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}$ ,  $(a'_i)^{(4)}$ ,  $(b_i)^{(4)}$ ,  $(b'_i)^{(4)}$ ,  $(p_i)^{(4)}$ ,  $(r_i)^{(4)}$ ,  $i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1.$$

Where we suppose,

- (O)  $(a_i)^{(5)}$ ,  $(a'_i)^{(5)}$ ,  $(a''_i)^{(5)}$ ,  $(b_i)^{(5)}$ ,  $(b'_i)^{(5)}$ ,  $(b''_i)^{(5)} > 0$ ,  $i, j = 28, 29, 30$   
 (P) The functions  $(a''_i)^{(5)}$ ,  $(b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}$ ,  $(r_i)^{(5)}$ :

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}(G_{31}, t) \leq (r_i)^{(5)} \leq (\hat{B}_{28})^{(5)}.$$

(Q)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$

**Definition of**  $(\hat{A}_{28})^{(5)}$ ,  $(\hat{B}_{28})^{(5)}$ :

Where  $(\hat{A}_{28})^{(5)}$ ,  $(\hat{B}_{28})^{(5)}$ ,  $(p_i)^{(5)}$ ,  $(r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$ .

They satisfy Lipschitz condition:

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}(G'_{31}, t) - (b''_i)^{(5)}(G_{31}, t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G'_{31})| e^{-(\hat{M}_{28})^{(5)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a''_i)^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 5$  then the function  $(a''_i)^{(5)}(T_{29}, t)$ , the **FIFTH augmentation coefficient** attributable would be absolutely continuous. .

**Definition of**  $(\hat{M}_{28})^{(5)}$ ,  $(\hat{k}_{28})^{(5)}$ :

- (R)  $(\hat{M}_{28})^{(5)}$ ,  $(\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1.$$

**Definition of**  $(\hat{P}_{28})^{(5)}$ ,  $(\hat{Q}_{28})^{(5)}$ :

- (S) There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}$ ,  $(\hat{k}_{28})^{(5)}$ ,  $(\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}$ ,  $(a'_i)^{(5)}$ ,  $(b_i)^{(5)}$ ,  $(b'_i)^{(5)}$ ,  $(p_i)^{(5)}$ ,  $(r_i)^{(5)}$ ,  $i = 28, 29, 30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1.$$

Where we suppose.

$(a_i)^{(6)}$ ,  $(a'_i)^{(6)}$ ,  $(a''_i)^{(6)}$ ,  $(b_i)^{(6)}$ ,  $(b'_i)^{(6)}$ ,  $(b''_i)^{(6)} > 0$ ,  $i, j = 32, 33, 34$

- (T) The functions  $(a''_i)^{(6)}$ ,  $(b''_i)^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}$ ,  $(r_i)^{(6)}$ :

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}(G_{35}, t) \leq (r_i)^{(6)} \leq (\hat{B}_{32})^{(6)}.$$

(U)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)}$   
 $\lim_{G \rightarrow \infty} (b''_i)^{(6)}(G_{35}, t) = (r_i)^{(6)}$

**Definition of**  $(\hat{A}_{32})^{(6)}$ ,  $(\hat{B}_{32})^{(6)}$ :

Where  $(\hat{A}_{32})^{(6)}$ ,  $(\hat{B}_{32})^{(6)}$ ,  $(p_i)^{(6)}$ ,  $(r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$ .

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}(G'_{35}, t) - (b''_i)^{(6)}(G_{35}, t)| < (\hat{k}_{32})^{(6)} |(G_{35}) - (G'_{35})| e^{-(\hat{M}_{32})^{(6)}t}.$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(6)}(T'_{33}, t)$  and  $(a''_i)^{(6)}(T_{33}, t)$ .  $(T'_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a''_i)^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 6$  then the function  $(a''_i)^{(6)}(T_{33}, t)$ , the **SIXTH augmentation coefficient** would be absolutely continuous. .

**Definition of**  $(\hat{M}_{32})^{(6)}$ ,  $(\hat{k}_{32})^{(6)}$ :

- $(\hat{M}_{32})^{(6)}$ ,  $(\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1.$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  :

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1 .$$

**Theorem 1:** if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad \boxed{T_i(0) = T_i^0 > 0} .$$

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} , \quad T_i(0) = T_i^0 > 0 .$$

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad T_i(0) = T_i^0 > 0 .$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0} .$$

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0} .$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, .$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} .$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} .$$

By

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t [(a_{13})^{(1)} G_{14}(s_{13}) - ((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{13}), s_{13})) G_{13}(s_{13})] ds_{13} .$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t [(a_{14})^{(1)} G_{13}(s_{13}) - ((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{13}), s_{13})) G_{14}(s_{13})] ds_{13} .$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t [(a_{15})^{(1)} G_{14}(s_{13}) - ((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{13}), s_{13})) G_{15}(s_{13})] ds_{13} .$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t [(b_{13})^{(1)} T_{14}(s_{13}) - ((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{13}), s_{13})) T_{13}(s_{13})] ds_{13} .$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t [(b_{14})^{(1)} T_{13}(s_{13}) - ((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{13}), s_{13})) T_{14}(s_{13})] ds_{13} .$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t [(b_{15})^{(1)} T_{14}(s_{13}) - ((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{13}), s_{13})) T_{15}(s_{13})] ds_{13}$$

Where  $s_{13}$  is the integrand that is integrated over an interval  $(0, t)$ .

**Proof:**

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)}, .$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} .$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} .$$

By

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t [(a_{16})^{(2)} G_{17}(s_{16}) - ((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{16}), s_{16})) G_{16}(s_{16})] ds_{16} .$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t [(a_{17})^{(2)} G_{16}(s_{16}) - ((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{16}), s_{17})) G_{17}(s_{16})] ds_{16} .$$

$$\begin{aligned} \bar{G}_{18}(t) &= G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)} \cdot \\ \bar{T}_{16}(t) &= T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)} \cdot \\ \bar{T}_{17}(t) &= T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)} \cdot \\ \bar{T}_{18}(t) &= T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)} (G(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$ .

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(M_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(M_{20})^{(3)}t}$$

By

$$\begin{aligned} \bar{G}_{20}(t) &= G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20}^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)} \cdot \\ \bar{G}_{21}(t) &= G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)} \cdot \\ \bar{G}_{22}(t) &= G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)} \cdot \\ \bar{T}_{20}(t) &= T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)} \cdot \\ \bar{T}_{21}(t) &= T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)} \cdot \\ \bar{T}_{22}(t) &= T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$ .

Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(M_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(M_{24})^{(4)}t}$$

By

$$\begin{aligned} \bar{G}_{24}(t) &= G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24}^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)} \cdot \\ \bar{G}_{25}(t) &= G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)} \cdot \\ \bar{G}_{26}(t) &= G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)} \cdot \\ \bar{T}_{24}(t) &= T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)} \cdot \\ \bar{T}_{25}(t) &= T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)} \cdot \\ \bar{T}_{26}(t) &= T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$ .

Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(M_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(M_{28})^{(5)}t}$$

By

$$\begin{aligned} \bar{G}_{28}(t) &= G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28}^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)} \cdot \\ \bar{G}_{29}(t) &= G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)} \cdot \\ \bar{G}_{30}(t) &= G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)} \cdot \\ \bar{T}_{28}(t) &= T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)} \cdot \\ \bar{T}_{29}(t) &= T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)} \cdot \\ \bar{T}_{30}(t) &= T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$ .

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} .$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} .$$

By

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32} \right)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) G_{32}(s_{(32)}) \right] ds_{(32)} .$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)} .$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)} .$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)} .$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)} .$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$ ..

(a) The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} \left( G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left( 1 + (a_{13})^{(1)}t \right) G_{14}^0 + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left( e^{(\hat{M}_{13})^{(1)}t} - 1 \right) .$$

From which it follows that

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)}t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1.

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$ .

The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that.

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} \left( G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}s_{(16)}} \right) \right] ds_{(16)} = \left( 1 + (a_{16})^{(2)}t \right) G_{17}^0 + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left( e^{(\hat{M}_{16})^{(2)}t} - 1 \right) .$$

From which it follows that

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)}t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[ \left( (\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$ .

(a) The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} \left( G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}s_{(20)}} \right) \right] ds_{(20)} =$$

$$\left( 1 + (a_{20})^{(3)}t \right) G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left( e^{(\hat{M}_{20})^{(3)}t} - 1 \right) .$$

From which it follows that

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[ \left( (\hat{P}_{20})^{(3)} + G_{21}^0 \right) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$ .

(b) The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}s_{(24)}} \right) \right] ds_{(24)} =$$

$$\left( 1 + (a_{24})^{(4)}t \right) G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left( e^{(\hat{M}_{24})^{(4)}t} - 1 \right) .$$

From which it follows that

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[ \left( (\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1.

(c) The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$\left( 1 + (a_{28})^{(5)}t \right) G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left( e^{(\hat{M}_{28})^{(5)}t} - 1 \right) .$$

From which it follows that

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ \left( (\hat{P}_{28})^{(5)} + G_{29}^0 \right) e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1.

(d) The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying GLOBAL EQUATIONS into itself .Indeed it is obvious that

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\mathcal{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left( 1 + (a_{32})^{(6)} t \right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\mathcal{M}_{32})^{(6)}} \left( e^{(\mathcal{M}_{32})^{(6)} t} - 1 \right).$$

From which it follows that

$$(G_{32}(t) - G_{32}^0) e^{-(\mathcal{M}_{32})^{(6)} t} \leq \frac{(a_{32})^{(6)}}{(\mathcal{M}_{32})^{(6)}} \left[ \left( (\hat{P}_{32})^{(6)} + G_{33}^0 \right) e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 6

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$ .

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\mathcal{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\mathcal{M}_{13})^{(1)}} < 1$  and to choose

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have.

$$\frac{(a_i)^{(1)}}{(\mathcal{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + \left( (\hat{P}_{13})^{(1)} + G_j^0 \right) e^{-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}} \right] \leq (\hat{P}_{13})^{(1)}.$$

$$\frac{(b_i)^{(1)}}{(\mathcal{M}_{13})^{(1)}} \left[ \left( (\hat{Q}_{13})^{(1)} + T_j^0 \right) e^{-\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)}.$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying GLOBAL EQUATIONS into itself.

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric

$$d \left( (G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\mathcal{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\mathcal{M}_{13})^{(1)} t} \right\}.$$

Indeed if we denote

**Definition of  $\tilde{G}, \tilde{T}$  :**

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\mathcal{M}_{13})^{(1)} s_{(13)}} e^{(\mathcal{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \left\{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\mathcal{M}_{13})^{(1)} s_{(13)}} e^{-(\mathcal{M}_{13})^{(1)} s_{(13)}} + \right.$$

$$(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\mathcal{M}_{13})^{(1)} s_{(13)}} e^{(\mathcal{M}_{13})^{(1)} s_{(13)}} +$$

$$\left. G_{13}^{(2)} |(a'_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\mathcal{M}_{13})^{(1)} s_{(13)}} e^{(\mathcal{M}_{13})^{(1)} s_{(13)}} \right\} ds_{(13)}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$|G^{(1)} - G^{(2)}| e^{-(\mathcal{M}_{13})^{(1)} t} \leq \frac{1}{(\mathcal{M}_{13})^{(1)}} \left( (a_{13})^{(1)} + (a'_{13})^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)} \right) d \left( (G^{(1)}, T^{(1)}); (G^{(2)}, T^{(2)}) \right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows.

**Remark 1:** The fact that we supposed  $(a'_{13})^{(1)}$  and  $(b'_{13})^{(1)}$  depending also on t can be considered as not conformal with the reality, however we have put this hypothesis in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{13})^{(1)} e^{(\mathcal{M}_{13})^{(1)} t}$  and  $(\hat{Q}_{13})^{(1)} e^{(\mathcal{M}_{13})^{(1)} t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}, i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \left\{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right\} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0.$$

**Definition of  $(\widehat{\mathcal{M}}_{13})^{(1)}_1, (\widehat{\mathcal{M}}_{13})^{(1)}_2$  and  $(\widehat{\mathcal{M}}_{13})^{(1)}_3$  :**

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$$G_{13} < (\widehat{\mathcal{M}}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{\mathcal{M}}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{\mathcal{M}}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{\mathcal{M}}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{\mathcal{M}}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{\mathcal{M}}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}, G_{15}$  and  $G_{13}, G_{14}$  respectively..

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below..



**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$  then  $T_{14} \rightarrow \infty$ .

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}.$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded. The same property}$$

holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions .

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$  and to choose

$(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  large to have.

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}.$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}.$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying .

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}.$$

Indeed if we denote

$$\text{Definition of } \widehat{G}_{19}, \widehat{T}_{19} : (\widehat{G}_{19}, \widehat{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19}).$$

It results

$$\begin{aligned} |\widehat{G}_{16}^{(1)} - \widehat{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a_{16}')^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a_{16}'')^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a_{16}'')^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)}. \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &\left( (a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{K}_{16})^{(2)} \right) d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right). \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows.

**Remark 1:** The fact that we supposed  $(a_{16}')^{(2)}$  and  $(b_{16}'')^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}, i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i'')^{(2)}t} > 0 \text{ for } t > 0.$$

**Definition of**  $(\widehat{M}_{16})^{(2)}_1, (\widehat{M}_{16})^{(2)}_2$  and  $(\widehat{M}_{16})^{(2)}_3$  :

**Remark 3:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$  . indeed if

$$G_{16} < (\widehat{M}_{16})^{(2)} \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively..

**Remark 4:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below..

**Remark 5:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$  then  $T_{17} \rightarrow \infty$ .

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}.$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results .}$$

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded. The same property}$$

holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} ((b_{18}'')^{(2)}((G_{19})(t), t)) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions .

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$  and to choose

$(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  large to have.

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ (\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left( \frac{(\bar{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)}.$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ ((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)}.$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself.

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric

$$d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}.$$

Indeed if we denote

$$\text{Definition of } \widehat{G}_{23}, \widehat{T}_{23} : ((\widehat{G}_{23}), (\widehat{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23})).$$

It results

$$\begin{aligned} |\widehat{G}_{20}^{(1)} - \widehat{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a_{20}')^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a_{20}'')^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a_{20}'')^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a_{20}'')^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$|G^{(1)} - G^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} \leq$$

$$\frac{1}{(\bar{M}_{20})^{(3)}} ((a_{20})^{(3)} + (a_{20}')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}) d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}); (G_{23})^{(2)}, (T_{23})^{(2)} \right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows.

**Remark 1:** The fact that we supposed  $(a_{20}'')^{(3)}$  and  $(b_{20}'')^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  and  $(\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}, i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a_i')^{(3)} - (a_i'')^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0.$$

**Definition of**  $(\bar{M}_{20})^{(3)}_1, (\bar{M}_{20})^{(3)}_2$  and  $(\bar{M}_{20})^{(3)}_3$  :

**Remark 3:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$  . indeed if

$$G_{20} < (\bar{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\bar{M}_{20})^{(3)})_1 - (a_{21}')^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\bar{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\bar{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$ ,  $G_{22}$  and  $G_{20}$ ,  $G_{21}$  respectively..

**Remark 4:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below..

**Remark 5:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(3)}((G_{23})(t), t)) = (b'_{21})^{(3)}$  then  $T_{21} \rightarrow \infty$ .

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21})^{(3)} - (b''_i)^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}.$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded. The same property}$$

holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b''_{22})^{(3)}((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions .

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$  and to choose

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have.

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)}.$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ ((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)}.$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying IN to itself.

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric

$$d((G_{27})^{(1)}, (T_{27})^{(1)}, (G_{27})^{(2)}, (T_{27})^{(2)}) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t} \}$$

Indeed if we denote

$$\text{Definition of } (\widehat{G}_{27}), (\widehat{T}_{27}) : ((\widehat{G}_{27}), (\widehat{T}_{27})) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$$

It results

$$\begin{aligned} |\widehat{G}_{24}^{(1)} - \widehat{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$|(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)}t} \leq$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) d(((G_{27})^{(1)}, (T_{27})^{(1)}); (G_{27})^{(2)}, (T_{27})^{(2)})$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows.

**Remark 1:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  and  $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}, i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0.$$

**Definition of**  $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$  and  $((\widehat{M}_{24})^{(4)})_3$  :

**Remark 3:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$  . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$  it follows  $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$  and by integrating  
 $G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively..

**Remark 4:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below..

**Remark 5:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$  then  $T_{25} \rightarrow \infty$ .

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b''_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)} :$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_4}$  By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded. The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities hold also for  $G_{29}$ ,  $G_{30}$ ,  $T_{28}$ ,  $T_{29}$ ,  $T_{30}$ :

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$  and to choose  $(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ (\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)} .$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ ((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)} .$$

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself.

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric

$$d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

$$\text{Definition of } (\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$$

It results

$$\begin{aligned} |\widehat{G}_{28}^{(1)} - \widehat{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} &\left( (a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)} \right) d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}); (G_{31})^{(2)}, (T_{31})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (35,35,36) the result follows.

**Remark 1:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}\} (T_{29}(s_{(28)}), s_{(28)}) ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)} t} > 0 \quad \text{for } t > 0.$$

**Definition of**  $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$  and  $((\widehat{M}_{28})^{(5)})_3$  :

**Remark 3:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$  . indeed if

$G_{28} < ((\widehat{M}_{28})^{(5)})$  it follows  $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)} G_{29}$  and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29}')^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30}')^{(5)} ((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$  ,  $G_{30}$  and  $G_{28}$  ,  $G_{29}$  respectively..

**Remark 4:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below..

**Remark 5:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(5)} ((G_{31})(t), t)) = (b_{29}')^{(5)}$  then  $T_{29} \rightarrow \infty$ .

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29}')^{(5)} - (b_i'')^{(5)} ((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)} :$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29}')^{(5)} (m)^{(5)} - \varepsilon_5 T_{29}$  which leads to

$$T_{29} \geq \left( \frac{(a_{29}')^{(5)} (m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \quad \text{If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29}')^{(5)} (m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \quad \text{By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded. The same property}$$

holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b_{30}'')^{(5)} ((G_{31})(t), t) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for  $G_{33}$  ,  $G_{34}$  ,  $T_{32}$  ,  $T_{33}$  ,  $T_{34}$  :

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$  and to choose

$(\widehat{P}_{32})^{(6)}$  and  $(\widehat{Q}_{32})^{(6)}$  large to have.

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ (\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} .$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ ((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} .$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  into itself.

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric

$$d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

$$\text{Definition of } (\widehat{G}_{35}), (\widehat{T}_{35}) : ((\widehat{G}_{35}), (\widehat{T}_{35})) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$$

It results

$$\begin{aligned} |\widehat{G}_{32}^{(1)} - \widehat{G}_{32}^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a_{32}')^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &(a_{32}'')^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + \\ &G_{32}^{(2)} |(a_{32}')^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a_{32}')^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows.

$$|(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} \leq$$

$$\frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a_{32}')^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}) d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}); (G_{35})^{(2)}, (T_{35})^{(2)} \right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows.

**Remark 1:** The fact that we supposed  $(a_{32}'')^{(6)}$  and  $(b_{32}'')^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$ ,  $i = 32, 33, 34$  depend only on  $T_{33}$  and respectively on  $(G_{35})(\text{and not on } t)$  and hypothesis can be replaced by a usual Lipschitz condition..

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 69 to 32 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0.$$

**Definition of**  $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$  and  $((\widehat{M}_{32})^{(6)})_3$  :

**Remark 3:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$ . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$  it follows  $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a_{33}')^{(6)}G_{33}$  and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33}')^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a_{33}')^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34}')^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a_{34}')^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$ ,  $G_{34}$  and  $G_{32}$ ,  $G_{33}$  respectively..

**Remark 4:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below..

**Remark 5:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$  then  $T_{33} \rightarrow \infty$ .

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33}')^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)} :$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33}')^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to

$$T_{33} \geq \left( \frac{(a_{33}')^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left( \frac{(a_{33}')^{(6)}(m)^{(6)}}{2} \right), t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded. The same property}$$

holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b_{34}')^{(6)}((G_{35})(t), t(t), t) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions.

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

- (a)  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying  
 $-(\sigma_2)^{(1)} \leq -(a_{13}')^{(1)} + (a_{14}')^{(1)} - (a_{13}'')^{(1)}(T_{14}, t) + (a_{14}'')^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$   
 $-(\tau_2)^{(1)} \leq -(b_{13}')^{(1)} + (b_{14}')^{(1)} - (b_{13}'')^{(1)}(G, t) - (b_{14}'')^{(1)}(G, t) \leq -(\tau_1)^{(1)}$ .

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$  :

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  :

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$  .

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :-

- (b) If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by  
 $(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$ , if  $(v_0)^{(1)} < (v_1)^{(1)}$   
 $(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}$ , if  $(v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}$ ,

and  $(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)} :$$

and analogously

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

and  $(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined respectively.

Then the solution satisfies the inequalities

$$G_{13}^0 e^{((s_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(s_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t} .$$

$$\left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[ e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \leq \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})} \left[ e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \right) .$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} .}$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} .$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[ e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t} .$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$ :-

Where  $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$   
 $(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$   
 $(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$   
 $(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)} .$

**Behavior of the solutions**

If we denote and define.

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  :

$\sigma_1^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying.  
 $-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} .$   
 $-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} .$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  :-

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots.

of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 .$

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and.

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  :-

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0 .$

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 .$

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :-

If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by.

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} .$   
 $(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$

and  $\boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}} .$

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} .$

and analogously

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$   
 $(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and  $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}} .$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} .$

Then the solution satisfies the inequalities

$G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)}t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t} .$

$(p_i)^{(2)}$  is defined.

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} .$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t} \right) .$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} .}$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} .$$

$$\frac{(b_{18})^{(2)}T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)}-(b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \frac{(a_{18})^{(2)}T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)}+(r_{16})^{(2)}+(R_2)^{(2)})} \left[ e^{((R_1)^{(2)}+(r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t} .$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :-

Where  $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$   
 $(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$   
 $(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$   
 $(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$ .

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

(a)  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying  
 $-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$   
 $-(\tau_2)^{(3)} \leq -(b_{20})^{(3)} + (b_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$ .

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

(b) By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$   
 and  $(b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$  and  
 By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$   
 and  $(b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$  :

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

(c) If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by  
 $(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$ , **if**  $(v_0)^{(3)} < (v_1)^{(3)}$   
 $(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$ , **if**  $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$ ,  
 and  $(v_0)^{(3)} = \frac{a_{20}^0}{a_{21}^0}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$ , **if**  $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$  ;  
 and analogously

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$ , **if**  $(u_0)^{(3)} < (u_1)^{(3)}$   
 $(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$ , **if**  $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$ , and  $(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$ , **if**  $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$  is defined .

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$  .

$\frac{(a_{22})^{(3)}G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})} \left[ e^{((S_1)^{(3)}-(p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \frac{(a_{22})^{(3)}G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)}-(a_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a_{22})^{(3)}t}$  .

$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t}$  .

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t}$  .

$\frac{(b_{22})^{(3)}T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq$

$\frac{(a_{22})^{(3)}T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[ e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$  .

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :-

Where  $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$   
 $(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$   
 $(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$   
 $(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$ .



**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

(d)  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) - (b''_{25})^{(4)}(G_{27}, t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  :

(e) By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  :

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the

$$\text{roots of the equations } (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

(f) If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } \boxed{(v_0)^{(4)} = \frac{\sigma_{24}^0}{\sigma_{25}^0}}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } \boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

are defined by 59 and 64 respectively

Then the solution satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined .

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a_{26}')^{(4)}t} + G_{26}^0 e^{-(a_{26}')^{(4)}t} \right]$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b_{26}')^{(4)}t} \right] + T_{26}^0 e^{-(b_{26}')^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a_{24}')^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b_{24}')^{(4)}$$

$$(R_2)^{(4)} = (b_{26}')^{(4)} - (r_{26})^{(4)}$$

### **Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

(g)  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a_{28}')^{(5)} + (a_{29}')^{(5)} - (a_{28}'')^{(5)}(T_{29}, t) + (a_{29}'')^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b_{28}')^{(5)} + (b_{29}')^{(5)} - (b_{28}'')^{(5)}((G_{31}), t) - (b_{29}'')^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  :

(h) By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  :

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the

$$\text{roots of the equations } (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

(i) If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

are defined respectively

Then the solution satisfies the inequalities

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined .

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t}$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \right) \leq G_{30}(t) \leq \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a_{30})^{(5)}t} + G_{30} e^{-(a_{30})^{(5)}t} \right]$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}}$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b_{30})^{(5)}t} \leq T_{30}(t) \leq$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :-

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions**

If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

(j)  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  :

(k) By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  :

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the

$$\text{roots of the equations } (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

(l) If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } \boxed{(v_0)^{(6)} = \frac{a_{32}^0}{a_{33}^0}}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } \boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

are defined respectively

Then the solution satisfies the inequalities

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined.

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t}$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \right) \leq G_{34}(t) \leq \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a_{34}')^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a_{34}')^{(6)}t} + G_{340} e^{-(a_{34}')^{(6)}t} \right]$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{33}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b_{34}')^{(6)}t} \right] + T_{34}^0 e^{-(b_{34}')^{(6)}t} \leq T_{34}(t) \leq$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$ :-

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a_{32}')^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b_{32}')^{(6)}$$

$$(R_2)^{(6)} = (b_{34}')^{(6)} - (r_{34})^{(6)}.$$

**Proof :** From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a_{13}')^{(1)} - (a_{14}')^{(1)} + (a_{13}'')^{(1)}(T_{14}, t) \right) - (a_{14}'')^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :-  $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$- \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

(a) For  $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

it follows  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$ .

In the same manner, we get

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

From which we deduce  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

(b) If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (\bar{C})^{(1)} (v_2)^{(1)} e^{[-(a_{14})^{(1)} ((v_1)^{(1)} - (v_2)^{(1)}) t]}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} ((v_1)^{(1)} - (v_2)^{(1)}) t]}} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}} \leq (\bar{v}_1)^{(1)} .$$

(c) If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$ , we obtain

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(1)}(t)$  :-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(1)}(t)$  :-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If  $(a_{13}'' )^{(1)} = (a_{14}'' )^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)} G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for the special case

Analogously if  $(b_{13}'' )^{(1)} = (b_{14}'' )^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

we obtain

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)} (T_{17}, t) \right) - (a''_{17})^{(2)} (T_{17}, t) v^{(2)} - (a_{17})^{(2)} v^{(2)} .$$

**Definition of**  $v^{(2)}$  :-  $\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$

It follows

$$- \left( (a_{17})^{(2)} (v^{(2)})^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left( (a_{17})^{(2)} (v^{(2)})^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)} \right) .$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

(d) For  $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$  .

In the same manner , we get

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ .

(e) If  $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$  we find like in the previous case,

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)} .$$

(f) If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$  , we obtain

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have .

**Definition of**  $v^{(2)}(t)$  :-

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}} .$$

In a completely analogous way, we obtain

**Definition of**  $u^{(2)}(t)$  :-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}} .$$

**Particular case :**

If  $(a_{16}'' )^{(2)} = (a_{17}'' )^{(2)}$  , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if  $(b_{16}'' )^{(2)} = (b_{17}'' )^{(2)}$  , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$ .

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t) \right) - (a_{21}'')^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)} .$$

**Definition of**  $v^{(3)}$  :-  $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right).$$

From which one obtains

(a) For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$ .

In the same manner, we get

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of**  $(\bar{v}_1)^{(3)}$  :-

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$ .

(b) If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case,

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

(c) If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$ , we obtain

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(3)}(t)$  :-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If  $(a_{20})^{(3)} = (a_{21})^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b_{20})^{(3)} = (b_{21})^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then



$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$ .

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :- 
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(v_1)^{(4)}, (v_0)^{(4)}$  :-

(d) For  $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

(e) If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)} .$$

(f) If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$  , we obtain

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)} , \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t) :-$

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{24})^{(4)} = (a''_{25})^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$  **this also defines  $(v_0)^{(4)}$  for the special case .**

Analogously if  $(b''_{24})^{(4)} = (b''_{25})^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then

$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of  $(u_0)^{(4)}$ .**

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)} :-$   $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)} :-$

(g) For  $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}, \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}, \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

(h) If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_2)^{(5)}]t}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}(\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}]t}} \leq (\bar{v}_1)^{(5)} .$$

(i) If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$ , we obtain

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (C)^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{28})^{(5)} = (a''_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then

$(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

we obtain

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of**  $v^{(6)}$  :-  $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

(j) For  $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (\bar{v}_1)^{(6)} .$$

(l) If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$  , we obtain

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)} , \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)} , \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{32}'' )^{(6)} = (a_{33}'' )^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case .**

Analogously if  $(b_{32}'' )^{(6)} = (b_{33}'' )^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

We can prove the following

**Theorem 3:** If  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  are independent on  $t$  , and the conditions

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0 ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined, then the system.

If  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  are independent on  $t$ , and the conditions .

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 .$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 .$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0 , .$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined are satisfied , then the system.

If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$ , and the conditions

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0 ,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined are satisfied , then the system.

If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$ , and the conditions

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0 ,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined are satisfied , then the system.

If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$ , and the conditions

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0 ,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined satisfied , then the system.

If  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  are independent on  $t$ , and the conditions

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined are satisfied , then the system.

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 .$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 .$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 .$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 .$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 .$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 .$$

has a unique positive solution , which is an equilibrium solution for the system.

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 .$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 .$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 .$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 .$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 .$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 .$$

has a unique positive solution , which is an equilibrium solution for .

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 .$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 .$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 .$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 .$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 .$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 .$$

has a unique positive solution , which is an equilibrium solution.

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0$$

has a unique positive solution , which is an equilibrium solution for the system

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0 .$$

(a) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0 .$$

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0 .$$

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0 .$$

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0 .$$

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0 .$$

**Definition and uniqueness of  $T_{14}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $T_{17}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first equations .

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

**Definition and uniqueness of  $T_{21}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(3)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $T_{25}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $T_{29}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $T_{33}^*$  :-**

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

(e) By the same argument, the equations 92,93 admit solutions  $G_{13}, G_{14}$  if

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - [(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$ .

(f) By the same argument, the equations 92,93 admit solutions  $G_{16}, G_{17}$  if

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - [(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0 .$$



Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi((G_{19})^*) = 0$ .

(g) By the same argument, the concatenated equations admit solutions  $G_{20}, G_{21}$  if

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$ .

(h) By the same argument, the equations of modules admit solutions  $G_{24}, G_{25}$  if

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$ .

(i) By the same argument, the equations (modules) admit solutions  $G_{28}, G_{29}$  if

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$ .

(j) By the same argument, the equations (modules) admit solutions  $G_{32}, G_{33}$  if

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G^*) = 0$ .

Finally we obtain the unique solution of 89 to 94

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution .

Finally we obtain the unique solution .

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0$  ,  $T_{17}^*$  given by  $f(T_{17}^*) = 0$  and.

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} .$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} .$$

Obviously, these values represent an equilibrium solution.

Finally we obtain the unique solution

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0$  ,  $T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution .

Finally we obtain the unique solution

$G_{25}^*$  given by  $\varphi(G_{27}) = 0$  ,  $T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]} .$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]}$$

Obviously, these values represent an equilibrium solution .

Finally we obtain the unique solution

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$  ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^*)]}$$

Obviously, these values represent an equilibrium solution.

Finally we obtain the unique solution

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$  ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]} .$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]}$$

Obviously, these values represent an equilibrium solution .

### ASYMPTOTIC STABILITY ANALYSIS

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  Belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij} .$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain .

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} .$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} .$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} .$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*\mathbb{G}_j .$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*\mathbb{G}_j .$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*\mathbb{G}_j .$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  Belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i .$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} .$$

taking into account equations (global)and neglecting the terms of power 2, we obtain .

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} .$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} .$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} .$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*\mathbb{G}_j .$$

$$\frac{d\mathbb{T}_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*\mathbb{G}_j .$$

$$\frac{d\mathbb{T}_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*\mathbb{G}_j .$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  Belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \quad \frac{\partial (b_i'')^{(3)}}{\partial G_j}(G_{23}^*) = s_{ij} .$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain .

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^* T_{21} .$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^* T_{21} .$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^* T_{21} .$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^* G_j .$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^* G_j .$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^* G_j .$$

If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote.

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i \quad , \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{25})''^{(4)}}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)} \quad , \quad \frac{\partial (b_i)''^{(4)}}{\partial G_j} ((G_{27})^*) = s_{ij} .$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain.

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^* T_{25} .$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^* T_{25} .$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^* T_{25} .$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^* G_j .$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^* G_j .$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^* G_j .$$

If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote.

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i \quad , \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{29})''^{(5)}}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)} \quad , \quad \frac{\partial (b_i)''^{(5)}}{\partial G_j} ((G_{31})^*) = s_{ij} .$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain.

$$\frac{dG_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^* T_{29} .$$

$$\frac{dG_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^* T_{29} .$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^* T_{29} .$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)}) T_{28}^* G_j .$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)}) T_{29}^* G_j .$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)}) T_{30}^* G_j .$$

If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Denote.

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i \quad , \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)} \quad , \quad \frac{\partial (b_i'')^{(6)}}{\partial G_j} ((G_{35})^*) = s_{ij} .$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain.

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^* T_{33} .$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^* T_{33} .$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^* T_{33} .$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)}) T_{32}^* G_j .$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)}) T_{33}^* G_j .$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)}) T_{34}^* G_j .$$

The characteristic equation of this system is

$$\begin{aligned} & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}) \{ ((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (q_{13})^{(1)} G_{13}^* \right] \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(14)} T_{14}^* + (b_{14})^{(1)} s_{(13),(14)} T_{14}^* \right) \\ & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}) (q_{13})^{(1)} G_{13}^* + (a_{13})^{(1)} (q_{14})^{(1)} G_{14}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(13)} T_{14}^* + (b_{14})^{(1)} s_{(13),(13)} T_{13}^* \right) \\ & \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & \left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ & + \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)} G_{15} \\ & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left( (a_{15})^{(1)} (q_{14})^{(1)} G_{14}^* + (a_{14})^{(1)} (a_{15})^{(1)} (q_{13})^{(1)} G_{13}^* \right) \\ & \left. \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}) s_{(14),(15)} T_{14}^* + (b_{14})^{(1)} s_{(13),(15)} T_{13}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & [((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^*]\} \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\
 & ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \\
 & ((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)}G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)}(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}G_{16}^*) \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(18)}T_{17}^* + (b_{17})^{(2)}s_{(16),(18)}T_{16}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & [((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)})(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(q_{20})^{(3)}G_{20}^*]\} \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(21)}T_{21}^* + (b_{21})^{(3)}s_{(20),(21)}T_{21}^* \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)})(q_{20})^{(3)}G_{20}^* + (a_{20})^{(3)}(q_{21})^{(1)}G_{21}^* \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(20)}T_{21}^* + (b_{21})^{(3)}s_{(20),(20)}T_{20}^* \\
 & ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \\
 & ((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \\
 & + ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} (q_{22})^{(3)}G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)}(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G_{20}^*) \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(22)}T_{21}^* + (b_{21})^{(3)}s_{(20),(22)}T_{20}^* \} = 0
 \end{aligned}$$

+

$$((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)})\{((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)})$$

$$\begin{aligned}
 & \left[ \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(25)} T_{25}^* + (b_{25})^{(4)} s_{(24),(25)} T_{25}^* \\
 & + \left( (\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)} \right) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(24)} T_{25}^* + (b_{25})^{(4)} s_{(24),(24)} T_{24}^* \\
 & \left( (\lambda)^{(4)} \right)^2 + \left( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} \\
 & \left( (\lambda)^{(4)} \right)^2 + \left( (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} \right) (\lambda)^{(4)} \\
 & + \left( (\lambda)^{(4)} \right)^2 + \left( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} \right) (\lambda)^{(4)} (q_{26})^{(4)} G_{26} \\
 & + \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) \left( (a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^* \right) \\
 & \left. \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & \left( (\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \left\{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \right\} \\
 & \left[ \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \\
 & + \left( (\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)} \right) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \\
 & \left( (\lambda)^{(5)} \right)^2 + \left( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} \\
 & \left( (\lambda)^{(5)} \right)^2 + \left( (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} \right) (\lambda)^{(5)} \\
 & + \left( (\lambda)^{(5)} \right)^2 + \left( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} \right) (\lambda)^{(5)} (q_{30})^{(5)} G_{30} \\
 & + \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) \left( (a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^* \right) \\
 & \left. \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & \left( (\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \left\{ (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \right\} \\
 & \left[ \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^*
 \end{aligned}$$

$$\begin{aligned}
 &+ \left( (\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)} \right) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 &\quad \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \\
 &\left( (\lambda)^{(6)} \right)^2 + \left( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} \\
 &\quad \left( (\lambda)^{(6)} \right)^2 + \left( (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} \right) (\lambda)^{(6)} \\
 &+ \left( (\lambda)^{(6)} \right)^2 + \left( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} \right) (\lambda)^{(6)} (q_{34})^{(6)} G_{34} \\
 &+ \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left( (a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \\
 &\left. \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right\} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

### Acknowledgments:

**The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's L:etters,Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidiation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive**

### REFERENCES

- [1] Dr K N Prasanna Kumar, Prof B S Kiranagi, Prof C S Bagewadi - MEASUREMENT DISTURBS EXPLANATION OF QUANTUM MECHANICAL STATES-A HIDDEN VARIABLE THEORY - published at: "International Journal of Scientific and Research Publications, Volume 2, Issue 5, May 2012 Edition".
- [2] DR K N PRASANNA KUMAR, PROF B S KIRANAGI and PROF C S BAGEWADI -CLASSIC 2 FLAVOUR COLOR SUPERCONDUCTIVITY AND ORDINARY NUCLEAR MATTER-A NEW PARADIGM STATEMENT - published at: "International Journal of Scientific and Research Publications, Volume 2, Issue 5, May 2012 Edition".
- [3] A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80th birthday.
- [4] FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188 .
- [5] HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford) [http://www.eolss.net.
- [6] MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", J. Geophys. Res., 111, D17204, doi:10.1029/2005JD006097.
- [7] STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" J. Atmos. Sci., 53, 980-1006.
- [8] FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" Nature, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010.

---

[9] A b c Einstein, A. (1905), "Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?", Annalen der Physik 18: 639 Bibcode 1905AnP...323..639E,DOI:10.1002/andp.19053231314. See also the English translation.



- [10] a b Paul Allen Tipler, Ralph A. Llewellyn (2003-01), Modern Physics, W. H. Freeman and Company, pp. 87–88, ISBN 0-7167-4345-0
- [11] a b Rainville, S. et al. World Year of Physics: A direct test of  $E=mc^2$ . Nature 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.
- [12] In F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy
- [13] Note that the relativistic mass, in contrast to the rest mass  $m_0$ , is not a relativistic invariant, and that the velocity is not a Minkowski four-vector, in contrast to the quantity  $dx^\mu$ , where  $dx^\mu$  is the differential of the proper time. However, the energy-momentum four-vector is a genuine Minkowski four-vector, and the intrinsic origin of the square-root in the definition of the relativistic mass is the distinction between  $dt$  and  $d\tau$ .
- [14] Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0
- [15] Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8
- [16] Hans, H. S.; Puri, S. P. (2003). Mechanics (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433
- [17] E. F. Taylor and J. A. Wheeler, Spacetime Physics, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.
- [18] Mould, Richard A. (2002). Basic relativity (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126
- [19] Chow, Tail L. (2006). Introduction to electromagnetic theory: a modern perspective. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392
- [20] [2] Cockcroft-Walton experiment
- [21] a b c Conversions used: 1956 International (Steam) Table (IT).
- [22] Assuming the dam is generating at its peak capacity of 6,809 MW.
- [23] Assuming a 90/10 alloy of Pt/Ir by weight, a  $C_p$  of 25.9 for Pt and 25.1 for Ir, a Pt-dominated average  $C_p$  of 25.8, 5.134 moles of metal, and 132 J.K<sup>-1</sup> for the prototype. A variation of  $\pm 1.5$  picograms is of course, much smaller than the actual uncertainty in the mass of the international prototype, which are  $\pm 2$  micrograms.
- [24] [3] Article on Earth rotation energy. Divided by  $c^2$ .
- [25] a b Earth's gravitational self-energy is  $4.6 \times 10^{-10}$  that of Earth's total mass, or 2.7 trillion metric tons. Citation: The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO), T. W. Murphy, Jr. et al. University of Washington, Dept. of Physics (132 kB PDF, here.).
- [26] There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be minimal coupling, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.
- [27] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", Physical Review D14:3432–3450 (1976).
- [28] A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", Physics Letters 59B:85 (1975).
- [29] F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", Physical Review D 30:2212.
- [30] Rubakov V. A. "Monopole Catalysis of Proton Decay", Reports on Progress in Physics 51:189–241 (1988).
- [31] S.W. Hawking "Black Holes Explosions?" Nature 248:30 (1974).
- [32] Einstein, A. (1905), "Zur Elektrodynamik bewegter Körper." (PDF), Annalen der Physik 17: 891–921, Bibcode 1905AnP...322...891E, DOI:10.1002/andp.19053221004. English translation.
- [33] See e.g. Lev B. Okun, The concept of Mass, Physics Today 42 (6), June 1969, p. 31–36, [http://www.physicstoday.org/vol-42/iss-6/vol42no6p31\\_36.pdf](http://www.physicstoday.org/vol-42/iss-6/vol42no6p31_36.pdf)
- [34] Max Jammer (1999), Concepts of mass in contemporary physics and philosophy, Princeton University Press, p. 51, ISBN 0-691-01017-X
- [35] Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", Foundations of Physics (Springer) 6: 115–124, Bibcode 1976FoPh....6..115E, DOI:10.1007/BF00708670
- [36] a b Janssen, M., Mecklenburg, M. (2007), From classical to relativistic mechanics: Electromagnetic models of the electron., in V. F. Hendricks, et al., , Interactions: Mathematics, Physics and Philosophy (Dordrecht: Springer): 65–134
- [37] a b Whittaker, E.T. (1951–1953), 2. Edition: A History of the theories of aether and electricity, vol. 1: The classical theories / vol. 2: The modern theories 1900–1926, London: Nelson
- [38] Miller, Arthur I. (1981), Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911), Reading: Addison–Wesley, ISBN 0-201-04679-2
- [39] a b Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), Séminaire Poincaré 1: 1–22

- [40] Philip Ball (Aug 23, 2011). "Did Einstein discover  $E = mc^2$ ?" Physics World.
- [41] Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* 42 (8): 540–543, DOI:10.1364/JOSA.42.000540
- [42] Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.
- [43] Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* 50 (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764
- [44] Ohanian, Hans (2008), "Did Einstein prove  $E=mc^2$ ", *Studies In History and Philosophy of Science Part B* 40 (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002
- [45] Hecht, Eugene (2011), "How Einstein confirmed  $E=mc^2$ ", *American Journal of Physics* 79 (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223
- [46] Rohrlich, Fritz (1990), "An elementary derivation of  $E=mc^2$ ", *American Journal of Physics* 58 (4): 348–349, Bibcode 1990AmJPh..58..348R, DOI:10.1119/1.16168
- [47] (1996). *Lise Meitner: A Life in Physics*. *California Studies in the History of Science*. 13. Berkeley: University of California Press. pp. 236–237. ISBN 0-520-20860-
- [48] R.C. Archibald (1914) *Time as a fourth dimension* *Bulletin of the American Mathematical Society* 20:409.
- [49] Gallier, Jean H. (2001). *Geometric methods and applications: for computer science and engineering*. Springer. p. 249. ISBN 0-387-95044-3., Chapter 8, page 249
- [50] Hermann Minkowski, "Raum und Zeit", 80. *Versammlung Deutscher Naturforscher (Köln, 1908)*. Published in *Physikalische Zeitschrift* 10 104–111 (1909) and *Jahresbericht der Deutschen Mathematiker-Vereinigung* 18 75–88 (1909). For an English translation, see Lorentz et al. (1952).
- [51] Einstein, Albert, 1926, "Space-Time," *Encyclopædia Britannica*, 13th ed.
- [52] Matolcsi, Tamás (1994). *Spacetime Without Reference Frames*. Budapest: Akadémiai Kiadó
- [53] Ellis, G. F. R.; Williams, Ruth M. (2000). *Flat and curved space-times* (2nd ed.). Oxford University Press. p. 9. ISBN 0-19-850657-0.
- [54] Petkov, Vesselin (2010). *Minkowski Spacetime: A Hundred Years Later*. Springer. p. 70. ISBN 90-481-3474-9., Section 3.4, p. 70
- [55] See "Quantum Spacetime and the Problem of Time in Quantum Gravity" by Leszek M. Sokolowski, where on this page he writes "Each of these hyper surfaces is spacelike, in the sense that every curve, which entirely lies on one of such hyper surfaces, is a spacelike curve." More commonly a space-like hyper surface is defined technically as a surface such that the normal vector at every point is time-like, but the definition above may be somewhat more intuitive.
- [56] Ehrenfest, Paul (1920). "How do the fundamental laws of physics make manifest that Space has 3 dimensions?" *Annalen der Physik* 61 (5): 440. Bibcode1920AnP...366...440E. DOI:10.1002/andp.19203660503. Also see Ehrenfest, P. (1917) "In what way does it become manifest in the fundamental laws of physics that space has three dimensions?" *Proceedings of the Amsterdam Academy* 20: 200.
- [57] Weyl, H. (1922) *Space, time, and matter*. Dover reprint: 284.
- [58] Tangherlini, F. R. (1963). "Atoms in Higher Dimensions". *Nuovo Cimento* 14 (27): 636.
- [59] Tegmark, Max (April 1997). "On the dimensionality of spacetime". *Classical and Quantum Gravity* 14 (4): L69–L75. arXiv:gr-c/9702052. Bibcode1997CQGra..14L..69T. DOI:10.1088/0264-9381/14/4/002. Retrieved 2006-12-16.
- [60] Dorling, J. (1970). "The Dimensionality of Time". *American Journal of Physics* 38(4): 539–40. Bibcode 1970AmJPh..38...539D. DOI:10.1119/1.1976386.
- [61] Barrow, J. D. (2002). *The Constants of Nature*. Pantheon Books. ISBN 0-375-42221-8.
- [62] Whitrow, G. J. (1955) "The Structure and Evolution of the Universe". *British Journal of the Philosophy of Science* 6: 13. Also see his (1959) *The Structure and Evolution of the Universe*. London: Hutchinson.
- [63] Jan Ambjørn, Jerzy Jurkiewicz, and Renate Loll - "The Self-Organizing Quantum Universe", *Scientific American*, July 2008
- [64] First Author: 1Mr. K. N.Prasanna Kumar has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding Author:drknpkumar@gmail.com
- [65] Second Author: 2Prof. B.S Kiranagi is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150

publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

- [66] Third Author: 3Prof. C.S. Bagewadi is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India