

Homotopy Perturbation Method for Solving Partial Differential Equations

R. Ashokan¹, M. Syed Ibrahim¹, L. Rajendran^{2,*}

¹Department of Mathematics, Madurai Kamaraj University, Madurai, Tamilnadu, India.

²Department of Mathematics, Sethu Institute of Technology, Kariapatti, Tamilnadu, India.

Corresponding Author: R. Ashokan

ABSTRACT: In this paper, we submitted a good tool to solve linear and nonlinear partial differential equations which is called homotopy perturbation method. This method makes hard problems so easy to solve, in our paper we gave some various examples for linear and nonlinear partial differential equations by using this method.

KEYWORDS: Homotopy perturbation method (HPM); Mathematical modeling; Partial differential equation.

Date of Submission: 30-03-2018

Date of acceptance: 14-04-2018

I. INTRODUCTION

Recently, many mathematicians seek new techniques to find exact or approximate solutions for nonlinear partial differential equations which describe different fields of science, physical phenomena, engineering, mechanics, and so on. Some modern methods have been appeared like homotopy perturbation method which is analytic technique for solving linear and nonlinear problems. The first mathematician proposed this method was Ji-Huan in 1999[1]. This method gives analytical exact and approximate solutions of nonlinear partial differential equations easily without transforming the equation or linearizing the problem with very good results. Some mathematician's author has used the homotopy perturbation method for solving partial differential problem [2-20]. In this work, we represent the solution for partial differential equations by homotopy perturbation method in linear and nonlinear type.

II. ANALYSIS OF HE'S HOMOTOPY PERTURBATION METHOD

To explain this method, we consider the following differential equation

$$D_o(u) - f(r) = 0, \quad r \in \Omega \tag{1}$$

with the boundary conditions

$$B_o(u, \frac{\partial u}{\partial n}) = 0, \quad r \in \Gamma \tag{2}$$

where D_o is a general differential operator, B_o is a boundary operator, $f(r)$ is a known analytical function and Γ is the boundary of the domain Ω . In general, the operator D_o can be divided into a linear part L and a non-linear part N . Eqn. (1) can therefore be written as

$$L(u) + N(u) - f(r) = 0 \tag{3}$$

By the homotopy technique, we construct a homotopy $v(r, p) : \Omega \times [0, 1] \rightarrow \Re$ that satisfies

$$H(v, p) = (1 - p)[L(v) - L(u_0)] + p[D_o(v) - f(r)] = 0. \tag{4}$$

$$H(v, p) = L(v) - L(u_0) + pL(u_0) + p[N(v) - f(r)] = 0. \tag{5}$$

where $p \in [0, 1]$ is an embedding parameter, and u_0 is an initial approximation of Eqn. (1) that satisfies the boundary conditions. From Eqns. (4) and (5), we have

$$H(v, 0) = L(v) - L(u_0) = 0 \tag{6}$$

$$H(v, 1) = D_o(v) - f(r) = 0 \tag{A7} \tag{7}$$

When $p=0$, Eqn. (4) and Eqn. (5) become linear equations. When $p=1$, they become non-linear equations. The process of changing p from zero to unity is that of $L(v) - L(u_0) = 0$ to $D_o(v) - f(r) = 0$. We first use the embedding parameter p as a “small parameter” and assume that the solutions of Eqns. (4) and (5) can be written as a power series in p :

$$v = v_0 + pv_1 + p^2v_2 + \dots \tag{8}$$

Setting $p = 1$ results in the approximate solution of Eqn. (1):

$$u = \lim_{p \rightarrow 1} v = v_0 + v_1 + v_2 + \dots \tag{9}$$

This series is convergent for most cases.

III. EXAMPLES

Example 1. Consider the following inhomogeneous heat equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \sin x \tag{10}$$

with initial condition

$$u(0, x) = 2 \cos x + \sin x \tag{11}$$

and a given solution

$$u(t, x) = 2 \cos x e^{-t} + \sin x. \tag{12}$$

To solve Eqn.(10) by Homotopy perturbation method, we will have

$$\frac{\partial u}{\partial t} - \frac{\partial u_0}{\partial t} = p \left[\frac{\partial^2 u}{\partial x^2} + \sin x \right] \tag{13}$$

Suppose that the solution of Eqn.(10) is the form

$$u = u_0 + pu_1 + p^2u_2 + \dots \tag{14}$$

Substituting Eqn.(14) into Eqn.(13) and equating the coefficients of like power p , we will have the set of differential equations:

$$\begin{aligned} p^0 : \frac{\partial u_0}{\partial t} - \frac{\partial u_0}{\partial t} &= 0 \\ p^1 : \frac{\partial u_1}{\partial t} &= \frac{\partial^2 u_0}{\partial x^2} + \sin x \\ p^2 : \frac{\partial u_2}{\partial t} &= \frac{\partial^2 u_1}{\partial x^2} \\ p^3 : \frac{\partial u_3}{\partial t} &= \frac{\partial^2 u_2}{\partial x^2} \\ p^4 : \frac{\partial u_4}{\partial t} &= \frac{\partial^2 u_3}{\partial x^2} \end{aligned} \tag{15}$$

and so on. Solve the system of Eqns.(14) to get the solutions

$$u_1 = -2 \cos x t$$

$$u_2 = 2 \cos x \frac{t^2}{2!} \tag{16}$$

$$u_3 = -2 \cos x \frac{t^3}{3!}$$

$$u_4 = 2 \cos x \frac{t^4}{4!}$$

Solution of Eqn. (10) will be derived by adding these terms, so

$$\begin{aligned} u(x, t) &= u_0 + u_1 + u_2 + \dots \dots \dots \\ &= 2 \cos x + \sin x - 2 \cos x t + 2 \cos x \frac{t^2}{2!} - 2 \cos x \frac{t^3}{3!} + 2 \cos x \frac{t^4}{4!} + \dots \\ &= 2 \cos x \left(1 - \frac{t}{1!} + \frac{t^2}{2!} - \dots \dots \dots \right) + \sin x \end{aligned}$$

$= 2 \cos x e^{-t} + \sin x$. (17) This is the exact solution (equ (12)) of the equ (10)

Example 2. Consider the following nonlinear PDEs

$$u_t = \frac{1}{3} u u_x - 3x \tag{18}$$

with initial condition

$$u(0, x) = 2 \tag{19}$$

and a given solution

$$u(t, x) = 2 \operatorname{sech} t - 3x \tanh t . \tag{20}$$

To solve Eqn.(16) by homotopy perturbation method, we will have

$$\frac{\partial u}{\partial t} - \frac{\partial u_0}{\partial t} = p \left[\frac{1}{3} u \frac{\partial u}{\partial x} - 3x \right] \tag{21}$$

Suppose that the solution of Eqn.(16) is the form

$$u = u_0 + p u_1 + p^2 u_2 + \dots \tag{22}$$

Substituting Eqn.(19) into Eqn.(16) and equating the coefficients of like power p , we will have the set of differential equations:

$$p^0 : \frac{\partial u_0}{\partial t} - \frac{\partial u_0}{\partial t} = 0$$

$$p^1 : \frac{\partial u_1}{\partial t} = \frac{u_0}{3} \frac{\partial u_0}{\partial x} - 3x$$

$$p^2 : \frac{\partial u_2}{\partial t} = \frac{u_0}{3} \frac{\partial u_1}{\partial x} + \frac{u_1}{3} \frac{\partial u_0}{\partial x}$$

(23)

$$p^3 : \frac{\partial u_3}{\partial t} = \frac{u_0}{3} \frac{\partial u_2}{\partial x} + \frac{u_1}{3} \frac{\partial u_1}{\partial x} + \frac{u_2}{3} \frac{\partial u_0}{\partial x}$$

$$p^4 : \frac{\partial u_4}{\partial t} = \frac{u_0}{3} \frac{\partial u_3}{\partial x} + \frac{u_1}{3} \frac{\partial u_2}{\partial x} + \frac{u_2}{3} \frac{\partial u_1}{\partial x} + \frac{u_3}{3} \frac{\partial u_0}{\partial x}$$

and so on. Solve the system of Eqns. (20) to get the solutions

$$u_1 = -3xt$$

$$u_2 = -t^2$$

$$u_3 = xt^3$$

$$u_4 = \frac{5t^4}{12}$$

and so on. Solution of Eqn. (16) will be derived by adding these terms, so

$$\begin{aligned} u(x, t) &= u_0 + u_1 + u_2 + \dots \dots \dots \dots \\ &= 2 - 3xt - t^2 + xt^3 + \frac{5}{12}t^4 + \dots \dots \dots \dots \\ &= 2 \left(1 - \frac{t^2}{2} + \frac{5t^4}{24} - \dots \dots \dots \right) - 3x \left(t - \frac{t^3}{3} + \dots \dots \dots \right) \\ u(t, x) &= 2 \operatorname{sech} t - 3x \tanh t. \end{aligned} \tag{25}$$

IV. CONCLUSION

In this paper, we used the homotopy perturbation method for solving some partial differential equations. We get the result is very effective and have an exact to find the solutions for the PDEs. Furthermore, HPM was successful implemented in approximating the solutions of nonlinear systems of PDEs.

REFERENCES

- [1]. J.H. He. Application of Homotopy Perturbation Method to Nonlinear Wave Equations. *Chaos, Solitons and Fractals*, 26(2000), 695-700.
- [2]. J.H. He. A coupling method of homotopy technique and perturbation technique for nonlinear problems. *Int. J. Nonlinear Mech.* 35(2000):37-43.
- [3]. J.H. He. Comparison of homotopy perturbation method and homotopy analysis method. *Appl. Math. Comput.* 156(2004):527-539.
- [4]. J. Biazar, H. Ghazvini. Exact solution for non-linear Schrodinger equations by Hes Homotopy perturbation method. *Physics Letters A* 366(2007):79-84.
- [5]. Zaid Odibat, Shaher Momani. A reliable treatment of homotopy perturbation method for Klein-Gordon equations. *Physics Letters A* 356(2007):351-357.
- [6]. Roba Al-Omary. Soliton solutions to systems of nonlinear partial differential equations using trigonometric-function method. Master Thesis. Jordan University of Science and Technology. 2011.
- [7]. S. Shukri, K. Al-Khaled. The extended tanh method for solving systems of nonlinear wave equations. *Appl. Math. Comput.* 217(2010): 1997-2006.
- [8]. N. H. Sweilam, M. M. Khader. Exact solutions of some coupled nonlinear partial differential equations using the homotopy perturbation method. *Computers and Mathematics with Applications*. 58 (2009):2134-2141.
- [9]. J. Biazar, K. Hosseini, P. Gholamin. Homotopy perturbation method for solving kdv and sawada-kotera equations. *Journal of Applied Mathematics*, 6(21)(2009):23-29.
- [10]. M. Alquran, M. Mohammad, Approximate solutions to system of nonlinear partial differential equations using homotopy perturbation method, *International Journal of Nonlinear Science* Vol.12(2011) No.4,pp.485-497.
- [11]. S. Loghambal, L. Rajendran, Mathematical modeling of diffusion and kinetics of amperometric immobilized enzyme electrodes, *Electrochimica Acta*, 55 (2010) 5230-5238.
- [12]. A. Meena, L. Rajendran, Analysis of pH - Based Potentiometric Biosensor using Homotopy perturbation method, *Chemical Engineering & Technology*, 33(2010) 1-10.
- [13]. A.Eswari, L.Rajendran, Analytical expressions of concentration and current in homogeneous catalytic reactions at spherical microelectrodes :Homotopy Perturbation approach, *Journal of Electroanalytical Chemistry*, 651 (2011) 173-184.
- [14]. S.Anitha, A.Subbiah, S.Subramaniam, L. Rajendran, Analytical solution of amperometric enzymatic reactions based on Homotopy perturbation method, *Electrochimica Acta*, 56 (2011) 3345-3352.
- [15]. S. Thiagarajan · A. Meena · S. Anitha, L. Rajendran, Analytical expression of the steady-state catalytic current of mediated bioelectrocatalysis and the application of He's Homotopy perturbation method, *JMathChem.* 6(2) 2011 96-104
- [16]. M. Subha, V. Ananthaswamy, L. Rajendran, A comment on Liao's Homotopy analysis method, *International Journal of Applied Sciences and Engineering Research*, Vol. 3, Issue 1, 2014,177-186.
- [17]. J.Saranya, L.Rajendran, L.Wang, C.Fernandez, A new mathematical modelling using homotopy perturbation method to solve nonlinear equations in enzymatic glucose fuelcells, *Chemical Physics Letters*, 662(2016),317-326.

- [18]. S. Pavithra, P. G. JansiRani, T. Iswarya and L. Rajendran, Enzymekineti c modelling and analytical solution of non linear rate equation in the transformation of d-methionineintol-methionine in batch reactor using the new homotopy perturbation method, *Quim. Nova*,39(10) (2016), 1184-1191.
- [19]. P.G. Jansi Rani, M. Kirthiga, Angela Molina, E.Laborda, L.Rajendran,Analytical solution of the convection-diffusion equation for uniformly accessible rotating disk electrodes via the homotopy perturbation method,*Journal of Electroanalytical Chemistry*, 799 (2017), 175-180.
- [20]. L. Rajendran, S. Anitha, Reply to “Comments on analytical solution of amperometric enzymatic Reaction sbasedon Homotopy perturbation method,Ji-Huan He, Lu-Feng Mo ElectrochimActa(2013), *ElectrochimicaActa*102(2013)474–476.

R. Ashokan." Homotopy Perturbation Method for Solving Partial Differential Equations." International Journal Of Modern Engineering Research (IJMER), vol. 08, no. 04, 2018, pp. 36–40.